

DEFORMATION QUANTIZATION OF SYMPLECTIC FIBRATIONS.

OLGA KRAVCHENKO

Abstract

A symplectic fibration is a fibre bundle in the symplectic category. We find the relation between deformation quantization of the base and the fibre, and the total space.

We use the weak coupling form of Guillemin, Lerman, Sternberg and find the characteristic class of deformation of symplectic fibration.

We also prove that the classical moment map could be quantized if there exists an equivariant connection.

Along the way we touch upon the general question of quantization with values in a bundle of algebras. We consider Fedosov's construction of deformation quantization in general. In Appendix we show how to calculate step by step the Fedosov connection, flat sections of the Weyl algebra bundle corresponding to functions and their star-product.

AMS classification: 58F06, 81S10, 58H15, 32G08

Key-words: deformation quantization, symplectic fibrations, moment map, Weyl algebra, Fedosov quantization.

Address: IRMA, Université Louis Pasteur, 7 rue René Descartes, 67084, Strasbourg, France

Email: ok@alum.mit.edu

1. INTRODUCTION: STATEMENT OF THE PROBLEM AND THE MAIN THEOREM.

Quantization is a map from functions on a (phase) space to operators on some Hilbert space. It involves a parameter (usually the Planck's constant h or $\hbar = \frac{h}{2\pi i}$). The product of two operators is given by some series in $\hbar$. The inverse of a quantization map allows one to get a noncommutative product on functions, namely, by taking the inverse of the product of corresponding operators. There is also the so called correspondence principle. The zero degree term in the $\hbar$ -decomposition of a noncommutative function (series in $\hbar$ with functional coefficients) should give a commutative product of functions. The term of degree one linearly depend on the Poisson bracket. This allows one to think of quantization as deformation of a structure of the algebra (Poisson algebra of functions on a manifold). In the formal deformation quantization one does not consider questions of convergence of series in $\hbar$. However, the formal deformation turns out to be a useful tool for describing global properties of a manifold. This concept of deformation quantization was described in [2]. When we say quantization we mean deformation quantization.

Fedosov found a beautiful construction of deformation quantization [11], [13] which we use here. Our article can be considered as an exercise in his domain, it is in fact a generalization of the first part of his article [12] and owes a lot to the methods of this article.

Quantization of a symplectic (or Poisson) manifold M is a construction of a noncommutative associative product on M . It is called a $*$ -product. $*$ -product is defined as a product on the algebra of series in a formal variable $\hbar$ with coefficients in $C^\infty(M)$. This noncommutative algebra $\mathbb{A}^\hbar$ should be a deformation of the algebra of functions on the manifold, $C^\infty(M)$. Let (M, ω_0) be a symplectic manifold. Then the symplectic form ω_0 defines a Lie-algebra structure on $C^\infty(M)$, called the Poisson structure. (The Jacobi identity follows from closedness of the form ω_0 .) For $f, g \in C^\infty(M)$ let $\{f, g\} = (df)^\sharp(g)$, where $\sharp : T^*M \rightarrow TM$, defined by ω_0 (see below section (2.2)).

Definition 1.1. Deformation quantization of a symplectic manifold (M, ω_0) is an associative algebra $(\mathbb{A}^\hbar, *)$ over $\mathbb{C}[[\hbar]]$ with an isomorphism $\epsilon : \mathbb{A}^\hbar / \hbar \mathbb{A}^\hbar \rightarrow C^\infty(M)$ s.t.

1. $\mathbb{A}^\hbar$ is flat as a $\mathbb{C}[[\hbar]]$ -module,
2. $\mathbb{A}^\hbar$ is separable and complete in $\hbar$ -topology,
3. for any $f, g \in C^\infty(M)$

$$\epsilon\left(\frac{i}{\hbar}(\check{f} * \check{g} - \check{g} * \check{f})\right) = \{f, g\}$$

for $\check{f}$ and $\check{g}$ s.t. $\epsilon\{\check{f}\} = f$ and $\epsilon\{\check{g}\} = g$,

4. The structure of the product $*$ on $\mathbb{A}^\hbar / \hbar^n \mathbb{A}^\hbar$ for all $n \geq 0$ is given by bidifferential operators.

DeWilde–Lecomte [8] and also Fedosov [11] proved that on any symplectic manifold there exists a quantization.

The following idea lies behind the Fedosov construction (see [9]): a Koszul-type resolution is considered for $C^\infty(M)[[\hbar]]$. Each term of the resolution has a noncommutative algebraic structure hence providing the algebra of functions with a new noncommutative product. Fedosov constructs such a resolution by using the differential forms on the manifold with values in the Weyl-algebra bundle. The main step then is to find a differential on it which respects the algebra structure. This differential is called Fedosov connection and is obtained by an iteration procedure from a torsion free symplectic connection on the manifold.

Lichnerowicz [23] showed that any connection on a symplectic manifold gives rise to a torsion-free symplectic connection. Hence one can get the Fedosov connection from any connection on a tangent bundle: first adding some tensor to make it symplectic (section 2.2) and then applying the iteration procedure.

We introduce a notion of an F -manifold:

Definition 1.2. An F -manifold Φ is the following triple (manifold, deformation of a symplectic form, a connection) :

$$\Phi = (M, \omega, \nabla),$$

where $\omega = \omega_0 + \hbar\alpha$, and $\alpha \in \Gamma(M, \Lambda^2 T^*M)[[\hbar]]$, a series in $\hbar$ with coefficients being closed 2-forms on the manifold.

In a recent article [16] a similar object is called a Fedosov manifold, namely, a symplectic manifold together with a symplectic connection. Indeed, these three objects (M, ω_0, ∇) define the first three terms in the $*$ -product:

- Classically, a manifold can be considered as an algebra of functions on it, that is M defines the structure of the commutative product.
- A symplectic form defines the Poisson structure and hence the term at $\hbar$.
- A connection defines the term at $\hbar^2$ (as follows from [23]).

It turns out that these three terms determine the higher terms. The deformation quantization theorem ([11]) can be stated as follows

Theorem 1.3. (*Fedosov*) An F -manifold (M, ω, ∇) uniquely determines a $*$ -product on the underlying manifold M .

Deligne [7] and Nest and Tsygan [28] showed that the class of isomorphisms of quantizations of a symplectic manifold M is determined by the class of the form ω in $H^2(M)[[\hbar]]$. It is called a *characteristic class of deformation*.

We study the deformation of the *twisted products* of two F -manifolds (B, ω_B, ∇^B) and (F, σ, ∇^F) . Our question is the following: how to define the product of two F -manifolds and what the $*$ -product on the total space is, that is what a “twisted product of quantizations” is.

We show under certain assumptions that a twisted product of two F -manifolds is again an F -manifold. So we want to relate the $*$ -products on these three manifolds.

One can regard the product $M = B \ltimes F$ as a fibre bundle $M \rightarrow B$ over a symplectic base B with a symplectic fibre F . Obviously, the product depends on how twisted the symplectic fibration $M = B \ltimes F$ is. This can be described by a connection on $M \rightarrow B$ which should be compatible with the symplectic form on M .

When G , the structure group of the bundle $M \rightarrow B$, acts by symplectomorphisms on fibres this bundle is a symplectic fibration (see [18], see below section 3). The total space M is symplectic and the fibres are symplectic submanifolds of it. We need the action to be Hamiltonian and G to be a finite dimensional Lie group.

The case of B being a cotangent bundle of some differentiable manifold X was first developed by Sternberg [30]. His construction describes the movement of a “classical particle” in Yang–Mills

field for any gauge group G and any differentiable X . Weinstein gave a general construction of a symplectic form on M in [32].

Here we are mainly interested in the particular case of the quantization with coefficients in the auxiliary bundle of fibrewise quantizations. This leads us to a more general case of quantization with coefficients in some bundle of algebras (some examples of such bundles are considered in [13]). Our case is somewhat different from the quantization with values in most other auxiliary bundle and is more difficult to perform. Fibres of the auxiliary bundle obtained from the symplectic fibration structure are noncommutative algebras. It is this noncommutativity of fibres which makes the quantization procedure more complicated.

However we believe that it is useful to understand its mechanism in order to see that quantization is a fundamental notion like some homology theory and hence it should respect a fibre bundle structure.

We make a new definition:

Definition 1.4. An F -bundle (with an underlying manifold B) is a triple

$$\Psi = (\Phi, A, \nabla^A)$$

where

- $\Phi = (B, \omega^B, \nabla^B)$ is an F -manifold,
- A is an auxiliary bundle of algebras over B ,
- ∇^A is a covariant derivative on A , which respects the algebraic structure on the fibres.

We construct an F -bundle from a symplectic fibration $M \rightarrow B$. Each fibre of $M \rightarrow B$ is an F -manifold, so we can quantize the fibres. If the connection ∇^F is G -invariant we can consider a new bundle A over the base B . This bundle A is a bundle of algebras of quantized functions on fibres. The fibre of A over a point b is the algebra of quantized functions on the fibre of $M \rightarrow B$ at the point b :

$$A_b = \mathbb{A}^{\hbar}(M_b). \quad (1)$$

The bundle A is defined by (1). The covariant derivative ∇^A on the bundle A is determined by the connection on the bundle $M \rightarrow B$. The construction of the covariant derivative is carried out in section (5.1) so that it respects the algebraic structure, that is it satisfies Leibnitz rule with respect to the product on A .

Such an F -bundle corresponds to an F -manifold modeled on M , the total space of the symplectic fibration, and hence provides a quantization of the total space. Our main theorem is:

Theorem 1.5. *Consider a symplectic fibration $M \rightarrow B$ with a standard fibre being an F -manifold (F, σ, ∇^F) and the base an F -manifold $\Phi = (B, \omega^B, \nabla^B)$. An F -bundle (Φ, A, ∇^A) , where $A_b = \mathbb{A}^{\hbar}(M_b)$ gives a quantization of the underlying manifold B with values in the auxiliary bundle. This also defines the quantization of the total space M .*

The main claim of this theorem is that quantization of the base with values in the auxiliary bundle (1) corresponds to a certain F -manifold $\Phi = (M, \omega, \nabla)$, where ω is a polynomial in $\hbar$ starting from a symplectic form on M .

To carry out the program first of all one has to construct a symplectic form on M . There is a one-parameter family of symplectic forms on the total space. The construction involves the notion of weak coupling limit of Guillemin, Lerman and Sternberg [18]. The behavior of the $*$ -product when this parameter tends to zero gives us a way to understand the relation between quantizations of the base and the fibre with the quantization of the total space.

Our main Theorem (1.5) can be reformulated as a statement about the solutions of two equations given in Theorem (5.6). We construct a twisted Weyl algebra bundle corresponding to F -bundle. Then we prove that

1. There exists r , a 1-form on the base with values in the twisted Weyl bundle, such that the initial connection becomes flat when one adds r to it.
2. For each section $\mathbf{a}$ of the auxiliary bundle there exists only one corresponding flat section of the twisted Weyl bundle.

Fedosov's quantization procedure is discussed in Section 2, calculations and examples are given in the Appendix. The classical setup for symplectic fibrations is discussed in Section 3 and the quantization of the moment map is presented in Section 4. Main results about the quantization of symplectic fibrations are given in Section 5, examples are discussed in the last section.

Notations. Repeated indices assume summation.

Grading and filtration of the Weyl algebra bundle are $\mathbb{Z}$ -grading and $\mathbb{Z}$ -filtration, we do not use the natural $\mathbb{Z}_2$ -grading on the differential forms.

Let $\mathcal{E}$ be a bundle over some manifold M . Then $\mathcal{A}^n(M, \mathcal{E})$ denotes C^∞ -sections of n -form bundle with values in the bundle $\mathcal{E}$,

$$\mathcal{A}^k(M, \mathcal{E}) = \Gamma(B, \Lambda^k T^*B \otimes \mathcal{E}).$$

$\mathcal{A}^n(M)$ denote the bundle of n -forms on M , and

$$\mathcal{A}(M, \mathcal{E}) = \oplus_{n=0}^{\infty} \mathcal{A}^n(M, \mathcal{E}).$$

The term “connection” is used in two senses, for a covariant derivative on any vector bundle, usually denoted by ∇ and also for a connection on a fibre bundle i.e. a splitting of the tangent bundle to the total space of a fibre bundle into a sum of a vertical and a horizontal subbundles.

Acknowledgments . I am deeply grateful to Ezra Getzler, Richard Melrose, Boris Tsygan and David Vogan for help and encouragement. My thanks also go to Alex Astashkevich, Paul Bressler, David Ellwood, Pavel Etingof, Daniel Grieser, Dima Kaledin, Eugene Lerman, and Andras Szenes for many fruitful discussions.

This work started at mathematics department at MIT and completed during my stay at IHES as a European postdoctoral fellow. It is a pleasure to acknowledge the support of these two institutions.

2. GENERALITIES ON DEFORMATION QUANTIZATION.

The subject of this section becomes nowadays fairly standard (see for example an excellent introduction to Fedosov quantization [22]).

2.1. Weyl algebra of a vector space.

Let E be a vector space with a non-degenerate skew-symmetric form ω . The algebra of polynomials on E is the algebra of symmetric powers of E^* , $S(E^*)$, and it has a skew-symmetric form on it which

is dual to ω . Let e be a point in E and $\{e^i\}$ denote its linear coordinates in E with respect to some fixed basis. Then $\{e^i\}$ define a basis in E^* . Let ω^{ij} be the matrix for the skew-symmetric form on E^* . Let us consider the power series in $\hbar$ with values in $S(E^*)$:

Definition 2.1. The Weyl algebra $W(E^*)$ of a vector space E^* is an associative algebra

$$W(E^*) = S(E^*)[[\hbar]] : \quad a(e, \hbar) = \sum_{k \geq 0} a_k(e) \hbar^k,$$

with the product structure given by the Moyal–Vey product:

$$a \circ b(e, \hbar) = \exp \left\{ -\frac{i\hbar}{2} \omega^{kl} \frac{\partial}{\partial x^k} \frac{\partial}{\partial z^l} \right\} a(x, \hbar) b(z, \hbar) \Big|_{x=z=e} \quad (2)$$

The Lie bracket is defined with respect to this product. We can look at this algebra as at a completion of the universal enveloping algebra of the Heisenberg algebra on $E^* \oplus \hbar\mathbb{C}$, namely, the algebra with relations

$$e^i \circ e^j - e^j \circ e^i = -i\hbar \omega^{ij}$$

where $\omega^{ij} = \omega(e^i, e^j)$ defines a Poisson bracket on E^* .

Hence one can define the Weyl algebra as

$$W = U(E^* \oplus \hbar\mathbb{C}).$$

Let us consider the product of the Weyl algebra and the exterior algebra of the space E^* : $W(E^*) \otimes \Lambda E^*$. Let dx^i be the basis in ΛE^* corresponding to e^i in $W(E^*)$.

There is a decreasing filtration on the Weyl algebra $W(E^*)$: $W_0 \supset W_1 \supset W_2 \supset \dots$ given by the degree of generators. e 's have degree 1 and $\hbar$ has degree 2:

$$W_p = \{\text{elements with degree} \geq p\}.$$

One can define a grading on W as follows

$$gr_i W = \{\text{elements with degree} = i\}.$$

it is isomorphic to W_i/W_{i+1} . One can see that the product (2) preserves the grading.

Definition 2.2. An operator on $W(E^*) \otimes \Lambda E^*$ is said to be of degree k if it maps $W_i \otimes \Lambda E^*$ to $W_{i+k} \otimes \Lambda E^*$ for all i .

Such an operator defines maps $gr_i W \otimes \Lambda E^*$ to $gr_{i+k} W \otimes \Lambda E^*$ for all i .

Definition 2.3. Derivation on $W(E^*) \otimes \Lambda E^*$ is a linear operator which satisfies the Leibnitz rule:

$$D(ab) = (Da)b + (-1)^{\tilde{a}\tilde{D}} a(Db)$$

where $\tilde{a}$ and $\tilde{D}$ are corresponding degrees. It turns out that all linear derivations are inner operators.

Lemma 2.4. Any linear derivation D on $W(E^*) \otimes \Lambda E^*$ is inner, i.e. namely there exists such $v \in W(E^*)$ so that $Da = \frac{i}{\hbar} [v, a]$ for any $a \in W(E^*)$

Proof. Indeed, $\frac{\partial}{\partial e^i} a = \frac{i}{2\hbar} [\omega_{ij} e^j, a]$.

So for any derivation one can get a formula: $Da = \frac{i}{\hbar} [\frac{1}{2} \omega_{ij} e^i D e^j, a]$. □

One can define two natural operators on the algebra $W(E^*) \otimes \Lambda E^*$: δ and δ^* of degree -1 and 1 correspondingly. δ is the lift of the “identity” operator

$$u : e^i \otimes 1 \rightarrow 1 \otimes dx^i$$

and δ^* is the lift of its inverse. On monomials $e^{i_1} \otimes \dots \otimes e^{i_m} \otimes dx^{j_1} \wedge \dots \wedge dx^{j_n} \in W^m(E^*) \otimes \Lambda^n E^*$ δ and δ^* can be written as follows:

$$\begin{aligned} \delta : e^{i_1} \otimes \dots \otimes e^{i_m} \otimes dx^{j_1} \wedge \dots \wedge dx^{j_n} &\mapsto \\ \sum_{k=1}^m e^{i_1} \otimes \dots \widehat{e^{i_k}} \dots \otimes e^{i_m} \otimes dx^{i_k} \wedge dx^{j_1} \wedge \dots \wedge dx^{j_n} \\ \delta^* : e^{i_1} \otimes \dots \otimes e^{i_m} \otimes dx^{j_1} \wedge \dots \wedge dx^{j_n} &\mapsto \\ \sum_{l=1}^n (-1)^l e^{j_l} \otimes e^{i_1} \otimes \dots \otimes e^{i_m} \otimes dx^{j_1} \wedge \dots \widehat{dx^{j_l}} \dots \wedge dx^{j_n}. \end{aligned}$$

Let a_0 be a projection of $a \in W(E^*) \otimes \Lambda E^*$ to $gr_0 W(E^*) \otimes \Lambda^0 E^*$, which is the center of the algebra, i.e. the summands in a which do not contain neither e -s or dx -s.

Lemma 2.5. *Operators δ and δ^* have the following properties:*

$$\delta a = dx^j \frac{\partial a}{\partial e^j} = \left[-\frac{i}{\hbar} \omega_{kl} e^k dx^l, a \right], \quad \delta^* a = y^j \iota_{\frac{\partial}{\partial x^j}} a, \quad \delta^2 = \delta^{*2} = 0$$

On monomials from $gr_m W(E^*) \otimes \Lambda^n E^*$

$$\delta \delta^* + \delta^* \delta = (m + n) Id,$$

where Id is the identity operator. Any element $a \in gr_m W(E^*) \otimes \Lambda^n E^*$ has a decomposition:

$$a = \frac{1}{m + n} (\delta \delta^* a + \delta^* \delta a) + a_0.$$

2.2. Symplectic connections (covariant derivatives).

The term *symplectic connection* in this section in fact must be changed to *symplectic covariant derivative* to avoid confusion with another symplectic connection notion in the next Chapter. However there is already an established practice to call a covariant derivative a connection which we decided to follow here. We hope that one can get used to distinguish one from the other from the context.

Let us consider connections on a manifold M .

Proposition 2.6. *Let ω be a skew-symmetric 2-form on $\mathcal{T}M$. Then ω must be closed in order for torsion-free connection ∇ preserving this form to exist.*

Proof. The skew-symmetry of ω is the following condition: $\omega(X, Y) = -\omega(Y, X)$. The connection ∇ is torsion-free when $\nabla_X Y - \nabla_Y X = [X, Y]$. Suppose such ∇ exists. Then it preserves the form ω when $\nabla \omega = 0$. This means that for all $X, Y, Z \in \mathcal{T}M$:

$$\nabla_X (\omega(Y, Z)) = \omega(\nabla_X Y, Z) + \omega(Y, \nabla_X Z) \quad (3)$$

Since $\omega(Y, Z)$ is a function $\nabla_X(\omega(Y, Z)) = X\omega(Y, Z)$. Then,

$$\begin{aligned}
X\omega(Y, Z) &= Y\omega(X, Z) + Z\omega(X, Y) \\
&= \omega(\nabla_X Y, Z) - \omega(\nabla_X Z, Y) - \omega(\nabla_Y X, Z) \\
&\quad + \omega(\nabla_Y Z, X) + \omega(\nabla_Z X, Y) - \omega(\nabla_Z Y, X) \\
&= \omega([X, Y], Z) - \omega([X, Z], Y) + \omega([Y, Z], X)
\end{aligned}$$

which is exactly the condition $d\omega = 0$. $\square$

Notice, that in the Riemannian case when the form is symmetric, there is a unique torsion free connection compatible with the form, the Levi-Civita connection. In the case of a skew-symmetric form there are plenty of connections compatible with the form, provided that the form is closed. So the statement of uniqueness of Levi-Civita connection in the Riemannian case is substituted by the requirement for the form to be closed in the skew-symmetric setting.

Here we are mostly interested in the case when M is a symplectic manifold, i.e. there is a symplectic form ω (a closed and nondegenerate 2-form on TM).

Definition 2.7. A connection which preserves a symplectic form is called a symplectic connection.

Any connection on a symplectic manifold gives rise to a symplectic connection:

Proposition 2.8. [23] [25]. *Let ω be a closed nondegenerate 2-form. Then for every connection ∇ there exists a three-tensor S , such that*

$$\tilde{\nabla} = \nabla + S$$

is a connection on TM compatible with ω .

Then for $X, Y \in TM$

$$\hat{\nabla}_X Y = \tilde{\nabla}_X Y - \frac{1}{2} \text{Tor}(X, Y)$$

defines a torsion-free connection compatible with the form ω . Here 2-form Tor is the torsion of $\tilde{\nabla}$

$$\text{Tor}(X, Y) = \tilde{\nabla}_X Y - \tilde{\nabla}_Y X - \tilde{\nabla}_{[X, Y]}$$

Then S is defined as follows:

$$S_X Y = \frac{1}{2} \{(\nabla_X \omega)(Y, \cdot)\}^\sharp,$$

where $\sharp : T^*M \rightarrow TM$ is the inverse to $\flat$, given by :

$$\flat : TM \rightarrow T^*M$$

$$u^\flat = \omega(u, \cdot) \text{ for } u \in TM.$$

Symplectic connections form an affine space with the associated vector space $\mathcal{A}^1(M, sp(2n))$, Lie algebra $sp(2n)$ valued one-forms on M .

2.3. Deformation quantization of a symplectic manifold.

Let M^{2n} be a symplectic manifold with a symplectic form ω . In local coordinates at a point x :

$$\omega = \omega_{ij} dx^i \wedge dx^j.$$

The symplectic form on a manifold M defines a Poisson bracket on functions on M . For any two functions $u, v \in C^\infty(M)$:

$$\{u, v\} = \omega^{ij} \frac{\partial u}{\partial x^i} \frac{\partial v}{\partial x^j} \quad (4)$$

where $(\omega^{ij}) = (\omega_{ij})^{-1}$.

We can define the bundle of Weyl algebras $\mathcal{W}_M$, with the fibre at a point $x \in M$ being the Weyl algebra of $\mathcal{T}_x^* M$. Let $\{e^1, \dots, e^{2n}\}$ be $2n$ generators in $\mathcal{T}_x^* M$, corresponding to dx^i . The form ω^{ij} defines pointwise Moyal–Vey product.

The filtration and the grading in $\mathcal{W}_M$ are inherited from $W(\mathcal{T}_x^* M)$ at each point $x \in M$. Denote by $\mathcal{W}^i$ the i -th graded component in $\mathcal{W}_M$:

$$\mathcal{W}_M = \oplus_i \mathcal{W}^i$$

A symplectic connection, ∇ , satisfying (3) can be naturally lifted to act on any symmetric power of the cotangent bundle (by the Leibnitz rule) and since the cotangent bundle $\mathcal{T}^* M \cong \mathcal{W}^1$ we can lift ∇ to be an operator on sections $\Gamma(M, \mathcal{W}^i)$ with values in $\Gamma(M, \mathcal{W}^i \otimes \mathcal{T}^* M)$. By abuse of notations this operator is also called ∇ .

It preserves the grading, in other words it is an operator of degree zero. It is clear that in general this connection is not flat: $\nabla^2 \neq 0$. Fedosov’s idea is that for $\mathcal{W}_M$ bundle one can add to the initial symplectic connection some operators not preserving the grading so that the sum gives a flat connection on the Weyl bundle.

Theorem 2.9. (Fedosov.) *There is a unique set of operators $r_k : \Gamma(M, \mathcal{W}^i) \rightarrow \Gamma(M, \mathcal{T}^* M \otimes \mathcal{W}^{i+k})$ such that*

$$D = -\delta + \nabla + r_1 + r_2 + \dots \quad (5)$$

is a flat connection and

$$\delta^* r_i = 0.$$

There is a one-to-one correspondence between formal series in $\hbar$ with coefficients in smooth functions $C^\infty(M)$ and horizontal sections of this connection:

$$Q : C^\infty(M)[[\hbar]] \rightarrow \Gamma_{\text{flat}}(M, \mathcal{W}_M). \quad (6)$$

Main idea of the proof is to use the following complex:

$$0 \rightarrow \Gamma(M, \mathcal{W}) \xrightarrow{\delta} \mathcal{A}^1(M, \mathcal{W}) \xrightarrow{\delta} \mathcal{A}^2(M, \mathcal{W}) \xrightarrow{\delta} \dots \quad (7)$$

This complex is exact since δ is homotopic to identity by δ^* . An equation for r_i for each $i > 1$ has the form

$$\delta(r_i) = \text{function}(\nabla, r_1, \dots, r_{i-1}). \quad (8)$$

However it is not difficult to show that this function is in the kernel of δ hence r_i exists.

First few steps in the construction of D , its flat sections and $*$ -product in coordinates are given in the Appendix.

The noncommutative structure on the Weyl bundle determines a $*$ -product on functions by this correspondence, namely for two functions $f, g \in \mathbb{C}^\infty(M)[[\hbar]]$;

$$f * g = Q^{-1}(Q(f) \circ Q(g)). \quad (9)$$

In fact, the equation $D^2 = 0$ is just the Maurer–Cartan equation for a flat connection. One can see the analogy with Kazhdan connection [15] on the algebra of formal vector fields. Notice that $\delta = dx^i \frac{\partial}{\partial e^i}$ is of degree -1 . The flatness of the connection is given by the recurrent procedure, namely starting from the terms of degree -1 and 0 one can get other terms step by step. While Kazhdan connection does not have a parameter involved it has the same structure – it starts with known -1 and 0 degree terms. Other terms are of higher degree and can be recovered one by one.

Let us also mention here that the connection D can be written as a sum of two terms – one is a derivation along the manifold, the usual differential d , and the other is an endomorphism of a fibre, let us call it Γ . Since all endomorphisms are inner one can write it as an adjoint action with respect to the Moyal product. Γ acts adjointly by an operator from $\Gamma(M, \mathcal{W})$ to $\Gamma(M, \mathcal{T}^*M \otimes \mathcal{W})$.

$$D = d + \Gamma = d + \frac{i}{\hbar}[\gamma, \cdot]_\circ, \quad (10)$$

where $\gamma \in \Gamma(M, \mathcal{T}^*M \otimes \mathcal{W})$. Then the equation $D^2 = 0$ becomes:

$$d\Gamma + \frac{1}{2}[\Gamma, \Gamma]_\circ = 0.$$

The same equation for γ then is as follows:

$$\omega + d\gamma + \frac{i}{\hbar} \frac{[\gamma, \gamma]_\circ}{2} = 0, \quad (11)$$

where ω is a central 2-form. This equation states that D^2 is given by an adjoint action of a central element, so it is zero. However it turns out to be very important which exactly form ω is given in the center by the connection D . Inner automorphisms of the Weyl algebra are given by the adjoint action by elements of the algebra (Lemma 2.4). Its central extension gives the whole algebra. Curvature of Fedosov connection is zero, however its lift to the central extension is nonzero and gives the characteristic class of quantization.

Definition 2.10. The characteristic class of the deformation is the cohomology class of the form $[\omega] \in H^2(M)[[\hbar]]$.

Quantization with values in a bundle of algebras. This subject was discussed at length in the book [12], but here we want to look at it from a slightly different angle. Given an F -manifold (B, ω, ∇^B) we know how to construct a map $C^\infty(B) \rightarrow \Gamma(B, \mathcal{W}_B)$. Let $\mathcal{L} \rightarrow B$ be a bundle of $\mathbb{C}[[\hbar]]$ -algebras. Now we want to generalize the problem of quantization and obtain a map:

$$Q^\mathcal{L} : \Gamma(B, \mathcal{L}) \rightarrow \Gamma(B, \mathcal{W}_B \otimes_{\mathbb{C}[[\hbar]]} \mathcal{L}).$$

In order to do that we need a connection $\nabla^\mathcal{L}$ on $\mathcal{L}$. Let $R^\mathcal{L} \in \mathcal{A}^2(B, \mathcal{L})$ be its curvature. Also the sections of $\mathcal{L}$ must commute with sections of the Weyl algebra bundle $\mathcal{W}_B$. Then we can define a connection on $\mathcal{W}_B \otimes_{\mathbb{C}[[\hbar]]} \mathcal{L}$ as the sum of connections.

$$\nabla = \nabla^B \otimes 1 + 1 \otimes \nabla^\mathcal{L}$$

with the curvature

$$R = R^B \otimes 1 + 1 \otimes R^\mathcal{L}.$$

We define a grading on $\mathcal{W}_B \otimes_{\mathbb{C}[[\hbar]]} \mathcal{L}$ as on $\mathcal{W}_B$. The operator $R^{\mathcal{L}}$ could have a degree if it changes the power of $\hbar$, and since the degree of $\hbar$ is 2 it could only be even: $R^{\mathcal{L}} = \sum R_{2k}^{\mathcal{L}}$, $k \leq 0$ where

$$R_{2k}^{\mathcal{L}} : \mathcal{A}^q(B, gr.(\mathcal{L})) \rightarrow \mathcal{A}^{q+2}(B, gr._{+2k}(\mathcal{L})).$$

One should add an extra term in the equation on r_i (8) for even $i = 2k$ one should add an extra term

$$\delta(r_{2k}) = \text{function}(\nabla, r_1, \dots, r_{2k-1}) + R_{2k}^{\mathcal{L}}.$$

Then like in Theorem 2.9 we can consider a complex similar to (7)

$$0 \rightarrow \Gamma(B, \mathcal{W}_B \otimes_{\mathbb{C}[[\hbar]]} \mathcal{L}) \xrightarrow{\delta} \mathcal{A}^1(B, \mathcal{W}_B \otimes_{\mathbb{C}[[\hbar]]} \mathcal{L}) \xrightarrow{\delta} \mathcal{A}^2(B, \mathcal{W}_B \otimes_{\mathbb{C}[[\hbar]]} \mathcal{L}) \xrightarrow{\delta} \dots$$

with δ acting only in $\mathcal{W}_B$. This complex is still exact (The $\otimes$ -product is over a field $\mathbb{C}[[\hbar]]$). Since $\delta R^{\mathcal{L}} = 0$ one can find a preimage of $R^{\mathcal{L}}$ and the reasoning is exactly as before. The flat connection and the corresponding flat sections are constructed similarly to the way it is described in Appendix.

However in the case when $\mathcal{L}$ has a Lie algebra structure and $R^{\mathcal{L}}$ is an inner action of the form $R^{\mathcal{L}} = \frac{i}{\hbar} \text{ad} H$ for some $H \in \mathcal{A}^2(B, \mathcal{L})$ the procedure changes! The adjoint action might start from the degree -2 term and one has to change not only the equations (8), but also the initial δ to balance it. This is exactly what happens in our case of symplectic fibrations and what makes it more interesting.

3. SYMPLECTIC FORMS ON SYMPLECTIC FIBRATIONS.

In this section we collect the facts known about symplectic fibrations: we give a definition and construct a one-parameter family of symplectic forms on the total space. One can find a nice exposition in the sixth chapter of the book [24], see also [18].

The meaning of the term *symplectic connection* used in this section is different from the definition 2.7. It is a connection which preserves the symplectic structure on fibres while in the previous Chapter 2 the symplectic connection was in fact a symplectic covariant derivative preserving a symplectic form on a symplectic manifold.

Symplectic fibrations.

Definition 3.1. A symplectic fibration is a locally trivial fibration $\pi : M \rightarrow B$ with a symplectic fibre (F, σ) whose structure group preserves the symplectic form σ on F . This means that

1. There is an open cover U_α of B and a collection of diffeomorphisms $\phi_\alpha : \pi^{-1}U_\alpha \rightarrow U_\alpha \times F$ such that the following diagram commutes:

$$\begin{array}{ccc} \pi^{-1}U_\alpha & \xrightarrow{\phi_\alpha} & U_\alpha \times F \\ & \searrow \pi & \swarrow pr \\ & U_\alpha & \end{array}$$

2. For the fibre over $b \in B$, $F_b = \pi^{-1}(b)$, let $\phi_\alpha(b)$ denote the restriction of ϕ_α to F_b followed by projection onto F , $\phi_\alpha(b) : F_b \rightarrow F$. Then

$$\phi_{\beta\alpha}(b) = \phi_\beta(b) \circ \phi_\alpha(b)^{-1} \in \text{Symp}(F, \sigma)$$

for all α, β and $b \in U_\alpha \cap U_\beta$.

If $\pi : M \rightarrow B$ is a symplectic fibration then each fibre F_b carries a symplectic structure $\sigma_b \in \Omega^2(F_b)$ defined by

$$\sigma_b = \phi_\alpha(b)^* \sigma$$

for $b \in U_\alpha$. The form is independent of α as follows from the definition. Also, if there is a G -invariant symplectic torsion-free connection ∇^F on F it defines a symplectic torsion-free connection ∇_b on each fibre F_b .

Definition 3.2. A symplectic form ω on the total space M of a symplectic fibration is called compatible with the fibration π if each fibre (F_b, σ_b) is a symplectic submanifold of (M, ω) , with σ_b being the restriction of ω to F_b .

Symplectic connections. Each symplectic form compatible with a symplectic fibration $\pi : M \rightarrow B$ defines a connection on it, i.e. a choice of splitting of the following short exact sequence of vector bundles:

$$0 \rightarrow \mathcal{V}M \rightarrow TM \rightarrow \pi^*TB \rightarrow 0.$$

Here $\mathcal{V}M$ is the canonically defined bundle of vertical tangent vectors, i.e. those fields which vanish on functions coming from the base. In other words the connection is the splitting of the tangent bundle into the sum

$$\Gamma : \quad TM = \mathcal{H}M \oplus \mathcal{V}M \tag{12}$$

such that $\pi^*TB = \mathcal{H}M$. The connection (12) is compatible with the symplectic form if at each point $x \in M$:

$$\mathcal{H}_x M := \{X \in \mathcal{T}_x M : \quad \Omega(X, V) = 0 \quad \text{for all } V \in \mathcal{V}_x M\}.$$

So each symplectic form whose restriction on fibres is nondegenerate defines a compatible connection. Namely, the horizontal subbundle consists of all vector fields which are perpendicular to the vertical ones with respect to the symplectic form.

Ingredients: a connection on a principal bundle and a Hamiltonian action along the fibres. Symplectic fibrations are associated fibre bundles to the principal bundles with a structure group being the group of symplectomorphisms of the fibre, so we have a principal G -bundle and a symplectic manifold (F, σ) to start with.

Let us first consider a principal G -bundle, i.e. a smooth manifold P with a smooth action $P \times G \rightarrow P$ which is free and transitive. Then the quotient $P/G = B$ is a manifold.

For the principal bundle a connection can be define by a so called connection 1-form. Namely, the fibres of a vertical subbundle $\mathcal{V}P$ are naturally identified with $\mathfrak{g}$ under the map: $\mathfrak{g} \rightarrow \text{Vect}(P)$ given by the infinitesimal action of G on P .

$$X \in \mathfrak{g} \mapsto \hat{X} \in \text{Vect}(P).$$

Hence the horizontal subbundle $\mathcal{H}P$ can be described not only as a kernel of the projection operator $Pr : \mathcal{T}P \rightarrow \mathcal{V}P$, but also as a kernel of a connection 1-form:

$$\lambda : \mathcal{T}P \rightarrow \mathfrak{g}.$$

It is a G -invariant form on the principal G -bundle P with values in the Lie algebra $\mathfrak{g}$, such that

$$\iota_{\hat{X}} \lambda = X, \quad \text{for } X \in \mathfrak{g}.$$

Now let G act on a symplectic manifold (F, σ) by symplectomorphisms, i.e. there is a group homomorphism

$$G \rightarrow \text{Symp}(F, \sigma) : g \mapsto \psi_g.$$

The infinitesimal action determines the Lie algebra homomorphism

$$\mathfrak{g} \rightarrow \text{Vect}(F, \sigma) : X \mapsto \hat{X}, \quad \text{defined by} \quad \hat{X} = \left. \frac{d}{dt} \right|_{t=0} \psi_{\exp(tX)} \quad (13)$$

A symplectic form determines a correspondence between functions and certain vector fields, called Hamiltonian vector fields:

$$C^\infty(F) \rightarrow \text{Vect}(F, \sigma) : H \mapsto X^H, \quad \text{defined by} \quad \iota_{X^H} \sigma = dH. \quad (14)$$

Definition 3.3. The action of G on F is called Hamiltonian if:

1. There is a lift $\mathfrak{g} \rightarrow C^\infty(F) : X \mapsto H_X$

$$\begin{array}{ccc} & & C^\infty(F) \\ & \nearrow & \downarrow \\ \mathfrak{g} & \longrightarrow & \text{Vect}(F, \sigma) \end{array}$$

This means that there is a Hamiltonian function H_X so that $\iota_{X^H} \sigma = dH_X$.

2. This map is a Lie algebra homomorphism:

$$H_{[X, Y]} = \{H_X, H_Y\}.$$

$[X, Y]$ is the Lie bracket in $\mathfrak{g}$ and $\{H_X, H_Y\}$ is the Poisson bracket in $C^\infty(F)$.

(If a group action satisfies only first condition it is called weakly Hamiltonian.) Let us also mention the following equality:

$$H_{[X, Y]} = \hat{X}H_Y - \hat{Y}H_X \quad (15)$$

Hamiltonian action determines a map $\mu : F \rightarrow \mathfrak{g}^*$, for each point $x \in F$ defined by

$$\langle \mu(x) | X \rangle = H_X(x)$$

for any $X \in \mathfrak{g}$, where $\langle \cdot | \cdot \rangle$ is the pairing: $\mathfrak{g}^* \times \mathfrak{g} \rightarrow \mathbb{C}$. This map is usually called a moment map, however for our purposes of quantization we will call a map $\mathfrak{g} \rightarrow C^\infty(F)$ also a moment map and it is this map of algebras which we are going to quantize.

Weak coupling: connection $\longleftrightarrow$ symplectic form. The following proposition in its present form is an adaptation for our purposes of a theorem about weak coupling form from [18].

Proposition 3.4. *Let $G \rightarrow \text{Symp}(F, \sigma) : g \mapsto \psi_g$ be a Hamiltonian action on (F, σ) with a moment map μ^F . Then every connection Γ on the principal G -bundle $P \rightarrow B$ over a symplectic manifold (B, ω^B) , gives rise to a one-parameter family of symplectic forms on the associated bundle $M = P \times_G F \rightarrow B$, which restricts to the forms σ_b on the fibres:*

$$\Omega_\epsilon = \epsilon^2 \omega^\Gamma + \pi^* \omega^B \quad (16)$$

where ϵ is a small parameter and ω^Γ is the coupling form, so that at a point $x \in M$, $\pi(x) = b$

$$\omega^\Gamma = \sigma_b + H_T$$

where $T \in \mathcal{A}^2(B)$: $T(X, Y) = -Pr([X^H, Y^H])$, $X, Y \in \mathcal{TB}$ and H_V is a Hamiltonian function of a vector field V with respect to the form σ_b , defined by μ_F .

Remark 3.5. Notice that σ_b is nonzero only on vertical vectors, while H_T is nonzero only on horizontal ones. This extra term H_T is needed for the form to be closed, ϵ makes the form Ω_ϵ to be nondegenerate.

Proof. Sketch (for the full proof see [18] or [24]). The main idea is to use the so called Weinstein universal phase space –

$$W = P \times \mathfrak{g}^*.$$

Given a connection on P the space W could be identified with the vertical subbundle of the cotangent bundle

$$W = P \times_G T^*G = \mathcal{V}^*P.$$

A connection is a splitting $\Gamma : \mathcal{TP} = \mathcal{HP} \oplus \mathcal{VP}$ and $\mathcal{V}^*P$ is defined as one-forms which vanish on horizontal vectors: $\mathcal{V}^*P = (\mathcal{HP})^\perp$. Hence it has a G -equivariant symplectic form coming from the canonical symplectic form on T^*P . Moreover, the action of the group G on W is Hamiltonian (see for example [1]).

The moment map $\mu^W : W \rightarrow \mathfrak{g}^*$ is given by the projection

$$pr_{\mathfrak{g}^*} : W \rightarrow \mathfrak{g}^*.$$

Then the symplectic reduction of $W \times F$ at 0 value of the moment map $\mu = \mu^W + \mu^F$ is exactly $M = P \times_G F$, and the symplectic form on M is inherited from W .

The explicit formula is obtained in the following way. Let the connection Γ be given by a connection 1-form, $\lambda_p : \mathcal{T}_p P \rightarrow \mathfrak{g}$. It determines a horizontal subbundle in $\mathcal{TP}$ by $\mathcal{H}_p P = \{v \in \mathcal{T}_p P \mid \lambda_p(v) = 0\}$.

$\mathcal{V}^*P = (\mathcal{HP})^\perp$ is also defined by λ . The connection $\lambda : \mathcal{TP} \rightarrow \mathfrak{g}$ and together with the action $\rho : \mathfrak{g} \rightarrow \mathcal{VP}$ define the injection

$$\iota_\lambda : \mathcal{V}^*P \hookrightarrow T^*P.$$

By definition of the connection 1-form, this injection is equivariant under the action of G and hence the 2-form

$$\omega_\lambda = \iota_\lambda^* \omega_{can} \in \mathcal{A}^2(\mathcal{V}^*P)$$

is invariant under the action of G . This pull-back of the canonical symplectic form on T^*P gives a closed 2-form on $\mathcal{V}^*P$.

The canonical 1-form α on T^*P is defined as follows. Let (p, s_p) be a point in T_p^*P , let also v be a tangent vector field in the tangent bundle $\pi : \mathcal{T}(T^*P) \rightarrow T^*P$ then at the point (p, s_p)

$$\langle \alpha | v \rangle_{(p, s_p)} = - \langle s_p | \pi_* v \rangle_p.$$

Then the pullback of the canonical one-form to $\mathcal{V}^*P$ is $\langle \iota_\lambda^* \alpha | v \rangle_{(p, s_p)} = - \langle pr_{\mathfrak{g}^*} s_p | \lambda(\pi_* v) \rangle_p$ so

$$\omega_\lambda = -d \langle pr_{\mathfrak{g}^*} | \lambda \rangle. \quad (17)$$

Now the form on $W \times F$ is $\omega_\lambda + \sigma$. We now want the form on the reduced space $M = (W \times F)/G$ at the regular value of the moment map $\mu^W + \mu^F = 0$. Since $\mu^W = pr_{\mathfrak{g}^*}$ the form on M becomes

$$\omega^\Gamma = d \langle \mu^F | \lambda \rangle + \sigma.$$

The one-form $\langle \mu^F | \lambda \rangle$ can be rewritten as the Hamiltonian H_λ . It should be understood in the following way. The connection 1-form $\lambda : \mathcal{TP} \rightarrow \mathfrak{g}$ defines the connection on the associated bundle $M = P \times_G F$. The horizontal subbundle in $\mathcal{TM}$ is the image of $\mathcal{HP}$ under the map $P \times F \rightarrow M$. By abuse of notation we call the map $\mathcal{TM} \rightarrow \mathfrak{g}$ also λ . From now on λ is a 1-form on M with values in the Lie algebra $\mathfrak{g}$. Hence H_λ is a 1-form on M .

Its differential gives a two form $d \langle \mu^F | \lambda \rangle = d(H_\lambda)$. Applied to two vectors $V, W \in \mathcal{TM}$:

$$d(H_\lambda)(V, W) = V H_\lambda(W) - W H_\lambda(V) - H_\lambda([V, W])$$

using (15) we get that it is nonzero only on horizontal vectors and gives the Hamiltonian of the curvature. We see also that ω^Γ restricted to fibres gives the symplectic form on the fibres. $\square$

This construction is quite general [18]: The symplectic fibrations with connection constructed this way turn out to include all symplectic fibre bundles with connection for which the holonomy group is a finite dimensional Lie group.

In local coordinates. Let us take a point $x \in M$. One can introduce a local frame $\{f_\alpha\}$ of vertical tangent bundle $\mathcal{VM}$ and a local frame $\{e_i\}$ in $\mathcal{TB}$ at a point $b = \pi(x)$ of B , with dual frames $\{f^\alpha\}$ and $\{e^i\}$. Using the connection we obtain a local frame on the tangent bundle $\mathcal{TM} = \pi^* \mathcal{TB} \oplus \mathcal{VM}$ at a point x .

Then the form can be written as a block matrix:

$$\Omega_\epsilon = \begin{vmatrix} \pi^* \omega^B + \epsilon^2 H_T & 0 \\ 0 & \epsilon^2 \sigma_b \end{vmatrix} \quad (18)$$

Hence the corresponding Poisson bracket is also a block matrix:

$$\begin{vmatrix} (\pi^* \omega^B + \epsilon^2 H_T)^{-1} & 0 \\ 0 & \epsilon^{-2} \sigma_b^{-1} \end{vmatrix}$$

We see that the Moyal product with respect to this form is a product of those on the base and on the fibres.

As for the connection, let $\Gamma_{\beta\gamma}^\delta$ be the Christoffel symbols of the symplectic connection on $\mathcal{TF}_b$ preserving the form σ_b along the fibre F_b . This connection can be written in the local coordinates as follows:

$$\nabla^F = d^F + \frac{i}{\hbar} [\Gamma_{\alpha\beta\gamma} f^\alpha f^\beta d\xi^\gamma, \cdot]. \quad (19)$$

where $\Gamma_{\alpha\beta\gamma} = (\sigma_b)_{\alpha\delta} \Gamma_{\beta\gamma}^\delta$.

4. QUANTUM MOMENT MAP.

Let (F, Σ, ∇) be an F -manifold. Here Σ is a deformation of the symplectic form σ :

$$\Sigma = \sigma + \hbar \sigma_1 + \hbar^2 \sigma_2 + \dots,$$

σ_i being closed 2-forms on F . Let $\mathbb{A}^\hbar(F)$ be the corresponding quantization of F with the characteristic class $[\Sigma] \in H^2(F)[[\hbar]]$. $\mathbb{A}^\hbar(F)$ is a noncommutative algebra of formal series in $\hbar$ with coefficients being smooth functions on F . The $*$ -product on $\mathbb{A}^\hbar(F)$ defines the Lie algebra structure:

$$[f, g]_* = \frac{i}{\hbar} (f * g - g * f), \quad \text{for } f, g \in \mathbb{A}^\hbar(F).$$

Let G be a group acting on F Hamiltonially (Definition 3.3). Let $\mathfrak{g} \rightarrow C^\infty(F)$ be its moment map. There is an induced action of G on $\mathbb{A}^\hbar(F)$. We want to quantize the moment map, namely, get a Lie algebra map from the algebra $\mathfrak{g}$ to the quantized algebra $\mathbb{A}^\hbar(F)$. However it is possible only up-to a two-cocycle in $C^\infty(F)[[\hbar]]$, so we get a projective representation or instead we should consider a central extension of $\mathbb{A}^\hbar(F)$. We could also slightly change a definition of a quantum moment map. We eliminate the central elements by considering the map into the adjoint representation of $\mathbb{A}^\hbar(F)$, the inner automorphisms $\text{Inn } \mathbb{A}^\hbar(F)$. They obviously inherit the Lie algebra structure from $\mathbb{A}^\hbar(F)$, so

Definition 4.1. A quantum moment map is a map of Lie algebras

$$\mu^\vee : \mathfrak{g} \rightarrow \text{Inn } \mathbb{A}^\hbar(F)$$

with the correspondence principle: $\lim_{\hbar \rightarrow 0} \mu^\vee(X)(f) = \{H_X, f\}$ for $X \in \mathfrak{g}$ and $f \in \mathbb{A}^\hbar(F)$.

Remark 4.2. This definition could be reformulated through an isomorphism of associative algebras: $\mu : U(\mathfrak{g}) \rightarrow \mathbb{A}^\hbar(F)$, such that on vector fields it gives:

$$\mu(X)^\vee(f) = [\mu(X), f]_*.$$

Proposition 4.3. Let G act Hamiltonially on (F, σ) with the Hamiltonian function H_X . Let

$$(F, \Sigma = \sigma + \sum_{i=1}^\infty \hbar^i \sigma_i, \nabla)$$

be an F -manifold such that ∇ is a G -invariant connection and assume that one can define functions $H_X^i, i = 1, 2, \dots$ as follows

$$\iota_{\hat{X}} \sigma_i = dH_X^i. \quad (20)$$

Let $\mathbb{A}^\hbar(F)$ be the algebra of quantized functions. Then the quantum moment map is given by

$$\mu(X) = H_X + \sum_{i=1}^\infty \hbar^i H_X^i \quad (21)$$

Also $\mu^\vee : \mathfrak{g} \rightarrow \text{Inn } \mathbb{A}^\hbar(F)$, $\mu(X)^\vee(f) = [\mu(X), f]_*$ is a homomorphism of Lie algebras:

$$\mu^\vee([X, Y]) = [\mu^\vee(X), \mu^\vee(Y)]_*. \quad (22)$$

Moreover, there are no higher terms in $\hbar$:

$$[\mu(X), f]_* = [H_X, f]_* = \{H_X, f\}. \quad (23)$$

*Proof.*¹ We are going to prove (21–23) by lifting the Hamiltonian H_X to a section of the Weyl algebra bundle, since the $*$ -product on $C^\infty(F)$ is defined through the the Weyl algebra bundle (6) and (9). Namely, let $\Gamma_{\text{D-flat}}(F, \mathcal{W}_F)$ be the space of flat sections Fedosov connection D constructed from ∇ corresponding to the quantization of F with the characteristic class Σ . Then there is a one-to-one correspondence

$$Q : C^\infty(F)[[\hbar]] \rightarrow \Gamma_{\text{D-flat}}(F, \mathcal{W}_F)$$

which defines a product on F

$$f * g = Q^{-1}(Q(f) \circ Q(g)).$$

¹I am grateful to Boris Tsygan for pointing out a gap in the proof in an earlier version of the paper.

The structure of a Lie algebra in Weyl algebra bundle $\mathcal{W}_F$ is defined by the fibrewise commutator

$$[\mathbf{a}, \mathbf{b}]_{\circ} = \mathbf{a} \circ \mathbf{b} - \mathbf{b} \circ \mathbf{a}, \quad \text{for } \mathbf{a}, \mathbf{b} \in \Gamma(F, \mathcal{W}_F)$$

Recall that the map $H : \mathfrak{g} \rightarrow C^{\infty}(F)$ is given by the condition

$$\iota_{\hat{X}} \sigma = dH_X, \quad (24)$$

where $\hat{X}$ is a field corresponding to X under the map $\mathfrak{g} \rightarrow \text{Vect}(F)$ (13). We find the image of a Hamiltonian H_X in $\Gamma_{\text{D-flat}}(F, \mathcal{W}_F)$ generalizing to the case of a deformed symplectic form the proof of Fedosov [13, Propositions 5.8.1,2],[14]. By construction the Fedosov connection D is G -equivariant since any element of the group G preserves the initial symplectic connection ∇ . An easy calculation shows also that flat sections of an equivariant connection D are also equivariant. It also means that the Lie derivative of D is zero with respect to any vector field $\hat{X}$:

$$[L_{\hat{X}}, D]_{\circ} = 0.$$

Since $L_{\hat{X}}$ and $D_{\hat{X}} = \iota_{\hat{X}} D + D \iota_{\hat{X}}$ are first order derivations commuting with D we can find an analogue of the Cartan homotopy formula for the Lie derivative on forms with values in the Weyl algebra bundle. The difference of $L_{\hat{X}}$ and $D_{\hat{X}}$ could only be an inner automorphism of the Weyl algebra bundle, we denote it $[Q(X), \cdot]_{\circ}$:

$$L_{\hat{X}} \mathbf{a} = (\iota_X D + D \iota_X) \mathbf{a} + [Q(X), \mathbf{a}]_{\circ}. \quad (25)$$

It is easy to see that the equality (25) is true in Darboux coordinates chosen so that the field $\hat{X}$ is just a pure derivation in the direction of only one of the coordinates. Then $D = D^0 = d + \delta$ and $Q(X) = Q^0(X) = \text{central section}(X, \hbar) - \iota_X \delta$. Let $D = D^0 + [\Delta \gamma, \cdot]_{\circ}$ be another flat connection, $\Delta \gamma$ being an equivariant one form in $\mathcal{W}_F$. Then since the $L_{\hat{X}}$ does not change and commutes with the new D as well, the right hand side of (25) must not change either, so one has to subtract $\iota_X \Delta \gamma$ from $Q^0(X)$.

We want to show that we could chose a *central section*($X, \hbar$) in $Q(X)$ to be equal to $\mu(X) = H_X + \hbar \cdots$ so that $Q(X) = \mu(X) - \iota_X \gamma$ becomes a quantization of the moment map, that is a flat section corresponding to H_X .

Recall that locally we can write a connection D as $Df = df + \frac{i}{\hbar} [\gamma, f]_{\circ}$ (10). The equation $D^2 = 0$ becomes (11):

$$\Sigma + d\gamma + \frac{i}{\hbar} \frac{[\gamma, \gamma]_{\circ}}{2} = 0. \quad (26)$$

Then since $L_{\hat{X}} D = 0$ from (25)

$$0 = \{\iota_{\hat{X}}(d + \frac{i}{\hbar} [\gamma, \cdot]_{\circ}) + (d + \frac{i}{\hbar} [\gamma, \cdot]_{\circ}) \iota_{\hat{X}}\}(D) - [\iota_X \gamma, D]_{\circ} = (\iota_{\hat{X}} d + d \iota_{\hat{X}})(D)$$

we get

$$(\iota_{\hat{X}} d + d \iota_{\hat{X}}) \gamma = 0. \quad (27)$$

Using (20),(24), and also (26) and (27) we find that indeed $Q(X)$ is a flat section of D :

$$\begin{aligned} DQ(X) &= d\{H_X + \hbar H_X^1 + \cdots\} - d(\iota_X \gamma) + \frac{i}{\hbar} [\gamma, \{H_X + \hbar H_X^1 + \cdots\}]_{\circ} - \frac{i}{\hbar} [\gamma, \iota_X \gamma]_{\circ} \\ &= \iota_X \Sigma + \iota_X d\gamma + 0 + \iota_X \left(\frac{i}{\hbar} \frac{[\gamma, \gamma]_{\circ}}{2} \right) = \iota_X \left(\Sigma + d\gamma + \frac{i}{\hbar} \frac{[\gamma, \gamma]_{\circ}}{2} \right) = 0. \end{aligned}$$

We also get that $\mu(X) = H_X + \Sigma_{i=1}^{\infty} \hbar^i H^i_X$, so the commutator

$$\begin{aligned} [\mu(X), f]_* &= \frac{i}{\hbar} Q^{-1} \{ [Q(X), Q(f)]_{\circ} \} = \frac{i}{\hbar} Q^{-1} \{ [(H_X + \hbar H_X^1 + \cdots - \iota_{\hat{X}} \gamma), Q(f)]_{\circ} \} \\ &= \frac{i}{\hbar} Q^{-1} \{ [-\iota_{\hat{X}} \gamma, Q(f)]_{\circ} \} = Q^{-1} \{ -\iota_{\hat{X}} (D - d) Q(f) \} = Q^{-1} \{ \iota_{\hat{X}} dQ(f) \} \\ &= L_{\hat{X}} f = \hat{X}(f) = \{H_X, f\}. \end{aligned}$$

Then since the action is Hamiltonian we get on the quantum level:

$$[\mu^{\vee}(X), \mu^{\vee}(Y)]_* = ad[H_X, H_Y]_* = ad(\{H_X, H_Y\}) = ad(H_{[X, Y]}) = \mu^{\vee}([X, Y]). \quad (28)$$

□

For the case $\Sigma = \sigma$, in the absence of additional terms $\hbar^i H^i$ we do not need to consider the adjoint representation. The notion of the quantum moment map repeats the classical one, it is a map from $\mathfrak{g}$ to $\mathbb{A}^{\hbar}(F)$:

Corollary 4.4. *Consider a quantization $\mathbb{A}^{\hbar}(F)$ obtained from the F -manifold (F, σ, ∇) . Let G act Hamiltonially on F with the Hamiltonian function H_X . Then the quantum moment map is a homomorphism of Lie algebras $\mu : \mathfrak{g} \rightarrow (\mathbb{A}^{\hbar}(F))$:*

$$\mu(X) = H_X \quad \text{and} \quad [\mu(X), f]_* = \{H_X, f\}.$$

Remark 4.5. We got that $\mu^{\vee} : \mathfrak{g} \rightarrow Inn \mathbb{A}^{\hbar}(F)$ is a map of Lie algebras. This gives the positive answer to a question posed in [33] that every classical moment map can be uniquely lifted to a quantum moment map to $\mathbb{A}^{\hbar}(F)$.

5. QUANTIZATION OF TWISTED PRODUCTS.

5.1. Auxiliary bundle: quantization of the fibres.

In this section we look at the bundle of quantizations along the fibres of our symplectic fibration $M \rightarrow B$ and construct a covariant derivative on this quantized bundle from the connection on the bundle $M \rightarrow B$.

Fibrewise quantization. Again, let $\pi : M \rightarrow B$ be a locally trivial fibration with a fibre F . Let F be equipped with a F -manifold (F, Σ, ∇) . Here we take Σ to be arbitrary characteristic class, i.e. cohomology class $[\Sigma] \in H^2(F)[[\hbar]]$.

$$\Sigma = \sigma + \hbar \sigma_1 + \hbar^2 \sigma_2 + \cdots,$$

where σ is a symplectic form on F .

Let $\mathbb{A}^{\hbar}(F)$ be the quantization of F , i.e. noncommutative algebra of formal series in $\hbar$ with coefficients being C^{∞} -functions on F . We are defining the bundle A over B such that its fibres are algebras of quantized functions on the fibres of the bundle $\pi : M \rightarrow B$, i.e. the fibre of A at a point $b \in B$ is

$$A_b = \mathbb{A}^{\hbar}(M_b).$$

The structure group G of $\pi : M \rightarrow B$ acts on F by symplectomorphisms hence there is a G -action on $\mathbb{A}^{\hbar}(F)$. Since $M = P \times_G F$ is an associated bundle to a G -bundle $P \rightarrow B$, this auxiliary bundle A is also associated to P with the fibre $\mathbb{A}^{\hbar}(F)$

$$A = P \times_G \mathbb{A}^{\hbar}(F).$$

Covariant derivative on the auxiliary bundle. If a connection on a fibre bundle is defined, there is a geometric way of putting a connection on the infinite dimensional bundle of smooth sections over fibres of a given finite dimensional fibre bundle.

In our case, covariant derivative on the bundle $A \rightarrow B$ respecting the algebra structure can be obtained from a connection 1-form on the principal bundle. Understanding of the formula for this covariant derivative is important for the sequel. We also find that the curvature of this covariant derivative equals to the adjoint action of the first summand in the coupling form.

In general, a choice of a connection on the principal G -bundle P determines a covariant derivative on any associated vector bundle (we follow an exposition in [3]).

Let $\lambda \in \mathcal{A}^1(P, \mathfrak{g})$ be a connection 1-form on a principal G -bundle P . Let also G act on some vector space E , and the action be given by the map ρ ,

$$\rho : G \rightarrow \text{End}(E).$$

Then the bundle $\mathcal{E} = P \times_G E \rightarrow B$ is an associated bundle to the principal bundle P . The space of differential forms on B with values in $\mathcal{E}$, $\mathcal{A}^k(B, P \times_G E)$, can be described as the subspace of the space of differential forms on P with values in E . This subspace is a space of all basic forms with values in E , $\mathcal{A}(P, E)_{bas}$. A *basic differential form* on a principal bundle P with a structure group G , taking values in the representation (E, ρ) of G , is an invariant and horizontal differential form, that is a form $\alpha \in \mathcal{A}(P, E)$ which satisfies

1. $g \cdot \alpha = \alpha, \quad g \in G$
2. $\iota(X)\alpha = 0$ for any vertical field X on P .

Lemma 5.1. *If $\alpha \in \mathcal{A}^q(P, E)_{bas}$, define $\alpha_B \in \mathcal{A}^q(B, P \times_G E)$ by*

$$\alpha_B(\pi_* X_1, \dots, \pi_* X_q)(b) = [p, \alpha(X_1, \dots, X_q)(p)],$$

where $p \in P$ is any point such that $\pi(p) = b$, and $X_i \in \mathcal{T}_p P$. Then α_B is well defined, and the map $\alpha \rightarrow \alpha_B$ is an isomorphism from $\mathcal{A}^q(P, E)_{bas}$ to $\mathcal{A}^q(B, P \times_G E)$.

As a particular case, there is a representation of the sections of $\mathcal{E}$ as G -equivariant functions on P with values in E . Let $C^\infty(P, E)^G$ denote the space of equivariant maps from P to E , that is those maps $s : P \rightarrow E$ that satisfy $s(p \cdot g) = \rho(g)s(p)$. There is a natural isomorphism between $\Gamma(B, P \times_G E)$ and $C^\infty(P, E)^G$, given by sending $s \in C^\infty(P, E)^G$ to s_B defined by

$$s_B(b) = [p, s(p)];$$

here p is any element of $\pi^{-1}(b)$ and $[p, s(p)]$ is the element of $\mathcal{E} = P \times_G E$ corresponding to $(p, s(p)) \in C^\infty(P, E)^G$. Infinitesimally, a function s in $C^\infty(P, E)^G$ satisfies the formula:

$$(X_P \cdot s)(b) + \rho(X)s(b) = 0, \quad \text{for } X \in \mathfrak{g},$$

where we also denote by ρ the differential of the representation ρ :

$$\rho : \mathfrak{g} \rightarrow \text{Vect}(E). \tag{29}$$

Given a connection 1-form λ on P one obtains the covariant derivative ∇ on the associated vector bundle $\mathcal{E}$ from the following commutative diagram:

$$\begin{array}{ccc} C^\infty(P, E)^G & \xrightarrow{d+\rho(\lambda)} & \mathcal{A}^1(P, E)_{bas} \\ \downarrow & & \downarrow \\ \Gamma(B, \mathcal{E}) & \xrightarrow{\nabla} & \mathcal{A}^1(B, \mathcal{E}) \end{array} \quad (30)$$

In other words, the covariant derivative can be written $(\nabla_X s)(b) = [p, (X^H s_P)(p)]$, where $X \in \mathcal{T}B$ and X^H is its horizontal lift to the principal bundle P . The formula for the covariant derivative on our auxiliary bundle below should be understood by means of the diagram (30).

Now let us return to our particular case, namely:

$$\mathcal{E} = A = P \times_G \mathbb{A}^{\hbar}(F), \text{ so } E = \mathbb{A}^{\hbar}(F),$$

Then (29) becomes a map $\rho : \mathfrak{g} \rightarrow Vect(\mathbb{A}^{\hbar}(F))$, or in other words it is given by the moment map $\mu^\vee : \mathfrak{g} \rightarrow Inn \mathbb{A}^{\hbar}(F)$.

Proposition 5.2. *Covariant derivative on $A \rightarrow B$ corresponding to a connection 1-form λ on $P \rightarrow B$ is given by the formula:*

$$\nabla^A f = df + [H_\lambda, f]_* = df + \{H_\lambda, f\}. \quad (31)$$

Its curvature is a 2-form on B with values in A :

$$R^A f = \{H_T, f\}, \quad \text{where } T(X, Y) = -Pr[X^H, Y^H], \quad Pr : \mathcal{T}P \rightarrow \mathfrak{g}.$$

Proof. The covariant derivative formula follows from (24) and the diagram (30). Then the general definition of the curvature of a covariant derivative $\nabla : \Gamma(B, E) \rightarrow \mathcal{A}^1(B, E)$ is

$$R(X, Y) = \nabla_{\tilde{X}} \nabla_{\tilde{Y}} - \nabla_{\tilde{Y}} \nabla_{\tilde{X}} - \nabla_{[\tilde{X}, \tilde{Y}]},$$

where $X, Y \in \mathcal{T}B$, and $\tilde{X}, \tilde{Y}$ are their horizontal lifts. In our particular case the expression for the curvature follows from the formulas (23) and (28). $\square$

5.2. Fedosov connection and flat sections on symplectic fibrations.

In this section we will show that the complex of differential forms with values in some twisted Weyl bundle gives a resolution of $\Gamma(B, A)$.

Bundle of sections. The structure of the bundle $\pi : M \rightarrow B$ is reflected in the representation of the space of functions on M as sections of a certain bundle over B . Namely, let $\mathcal{F}$ be the bundle over B , such that a fibre over $b \in B$ is the space of series in $\hbar$ with coefficients being functions on the fibre M_b of the bundle $\pi : M \rightarrow B$:

$$\mathcal{F}_b = C^\infty(M_b)[[\hbar]].$$

Then $C^\infty(M)[[\hbar]]$ can be represented as sections of the bundle $\mathcal{F}$:

$$C^\infty(M)[[\hbar]] = \Gamma(B, \mathcal{F}) \quad (32)$$

Hence, we can try to obtain a quantization of M by quantization of the bundle $\mathcal{F} \rightarrow B$. This leads us to consider the twisted Weyl algebra bundle over B .

Indeed, let $\mathcal{W}_M$ be the Weyl algebra bundle on M . Then consider $\mathcal{W}_{M/B}$, the bundle on B with the fibers $\mathcal{W}_b$ being the restriction of the bundle $\mathcal{W}_M$ to the fibre M_b , $b \in B$. Given a symplectic connection (12) on the bundle $\pi : M \rightarrow B$ we can talk about the following isomorphism on the level of sections

$$\beta : \Gamma(M, \mathcal{W}_M) \rightarrow \Gamma(B, \mathcal{W}_B \otimes_{\mathbb{C}[[\hbar]]} \Gamma(M/B, \mathcal{W}_{M/B})) \quad (33)$$

or more generally for differential forms:

Lemma 5.3. *Given a symplectic connection:*

$$\mathcal{T}M = \mathcal{H}M \oplus \mathcal{V}M, \quad \mathcal{H}_z M \cong \pi^* \mathcal{T}_{\pi(z)} B$$

there is an isomorphism:

$$\mathcal{A}^n(M, \mathcal{W}_M) \rightarrow \oplus_{p+q=n} \mathcal{A}^p(B, \mathcal{W}_B \otimes_{\mathbb{C}[[\hbar]]} \mathcal{A}^q(M/B, \mathcal{W}_{M/B})) \quad (34)$$

Proof. At each point z of M the Weyl algebra can be defined as the universal enveloping algebra of the Heisenberg algebra of $\mathcal{T}_z^* M \oplus \hbar \mathbb{R}$. Universal enveloping algebra is by definition a quotient of the tensor algebra by a certain ideal. The ideal is

$$I = \{e \otimes f - e \otimes f + i\hbar \omega^{-1}(e, f)\}, \quad \text{for } e, f \in \mathcal{T}_z^* M$$

and ω the symplectic form on $\mathcal{T}_z^* M$. The connection on $\mathcal{T}M$ splits the ideal into a sum of a horizontal and vertical ideals:

$$I = I_H + I_V.$$

Thanks to the splitting of the symplectic form (18) vertical ideal I_V is also an ideal in the tensor algebra of $\mathcal{V}^* M$, which leads to the result. $\square$

Remark 5.4. Notice that we were talking about the isomorphism of bundles, but we did not show yet that it is an isomorphism of algebras – it is the subject of the section 5.3.

We see that one can consider fiberwise quantization as a first step in quantization of the total space. This leads to the auxiliary bundle described in Section 5.1. The quantization map along the fibres:

$$Q^{\text{fibre}} : \Gamma(B, \mathcal{F}) \rightarrow \Gamma(B, A) \quad (35)$$

Twisted Weyl algebra bundle. Let $\mathcal{W}_B$ be the Weyl algebra bundle corresponding to the F -manifold $\{B, \omega, \nabla^B\}$. Consider the twisted bundle

$${}^A \mathcal{W}_B = \mathcal{W}_B \otimes_{\mathbb{C}[[\hbar]]} A \rightarrow B.$$

Thanks to the splitting of the form (18) the sections of $\mathcal{W}_B$ and sections of A commute with each other.

Remark 5.5. $\Gamma(B, {}^A \mathcal{W}_B)$ can be considered as a space of fiberwise flat sections of the bundle obtained by β (33): $\Gamma(B, \mathcal{W}_B \otimes_{\mathbb{C}[[\hbar]]} \Gamma(M/B, \mathcal{W}_{M/B}))$. Indeed, the F -manifold (F, σ, ∇^F) induces a structure of F -manifold on each fibre. It defines corresponding isomorphism of flat sections of Weyl algebra bundles:

$$A = \Gamma_{\text{flat}}(M/B, \mathcal{W}_{M/B}) \quad (36)$$

This bundle ${}^A\mathcal{W}_B$ is a bundle of graded algebras with degrees assigned as in the original $\mathcal{W}_B$ bundle, namely,

$$\deg(\hbar) = 2, \quad \deg(e^i) = 1, \quad e^i\text{'s being generators of } \mathcal{W}_B. \quad (37)$$

Let

$$({}^A\mathcal{W}_B)_n = \{s \in {}^A\mathcal{W}_B, \quad \text{such that} \quad \deg(s) \geq n\}.$$

Then let us also define the grading

$$gr_n({}^A\mathcal{W}_B) = \{s \in {}^A\mathcal{W}_B, \quad \text{such that} \quad \deg(s) = n\}.$$

It is isomorphic to $({}^A\mathcal{W}_B)_n / ({}^A\mathcal{W}_B)_{n+1}$. The pointwise noncommutative product on ${}^A\mathcal{W}_B$ is inherited from the Moyal–Vey product $\circ$ for $\mathcal{W}_B$ and the noncommutative product $*$ on the auxiliary bundle A . Let us denote the product on ${}^A\mathcal{W}_B$ also by $\circ$. The noncommutative product on A contains terms in different degrees in $\hbar$, but does not have any other degree bearing terms. Hence, the product on ${}^A\mathcal{W}_B$ is not preserving the grading anymore.

The Lie algebra structure is defined as well. For $f, g \in \Gamma(B, {}^A\mathcal{W}_B)$:

$$[f, g] = [f, g]_\circ + [f, g]_*.$$

The symplectic connection on B and the connection on the auxiliary bundle give rise to a connection on ${}^A\mathcal{W}_B$:

$$\nabla = \nabla^B \otimes 1 + 1 \otimes \nabla^A.$$

In what follows we will omit the tensor signs, this should not cause a confusion.

The curvature of this connection is

$$R = R^B + R^A$$

Following the general scheme we want to flatten the connection ∇ by adding operators of degree -1 and higher.

Main theorem.

Theorem 5.6. *The equation $(D^A)^2 = 0$ for the connection*

$$D^A : \mathcal{A}^q(B, {}^A\mathcal{W}_B) \rightarrow \mathcal{A}^{q+1}(B, {}^A\mathcal{W}_B)$$

has a solution in the form:

$$D^A = d + \frac{i}{\hbar}[\gamma, \cdot], \quad (38)$$

where $\gamma \in \text{Inn} {}^A\mathcal{W}_B$ is a sum of operators of degree ≥ -1 , so that $d + \frac{i}{\hbar}[\gamma, \cdot] = \nabla + \delta + r$, in particular $\deg(\delta) = -1$ and

$$r = \sum_{k \geq 1} r_k, \quad r_k \in gr_k {}^A\mathcal{W}_B.$$

When r satisfies the normalization condition $\delta^{-1}r = 0$ the solution is unique.

Flat sections of this connection are in one-to-one correspondence with sections of the auxiliary bundle, $\Gamma(B, A)$:

$$Q^A : \Gamma(B, A) \rightarrow \Gamma_{\text{flat}}(B, \mathcal{W}). \quad (39)$$

Proof. Main equation is $\omega + d\gamma + \frac{i}{\hbar}[\gamma, \gamma] = 0$. It can be rewritten

$$R + \frac{1}{2}[\nabla, \delta] + \frac{1}{2}[\nabla, r] + \delta^2 + \frac{1}{2}[\delta, r] + r^2 = 0.$$

For each $k > 1$ we get an equation expressing r_k in terms of $r_i, i \leq k$:

$$[\delta, r_k] = [\nabla, r_{k-1}] + \sum_{i=1}^{k-2} [r_i, r_{k-i}]. \quad (40)$$

However the equation on $r - 1$ which must kill the curvature gives an unusual term in -1 degree. Namely, the equation on r_1 : $\delta r_1 = R^B + R^A$ gives a solution which is a sum of two terms in degrees 1 and -1 (sic!). Indeed, in coordinates:

$$r_1 = \delta^{-1}(R^B + R^A) = \delta^{-1}(R_{ijkl}^B e^i e^j + R_{kl}^A) dx^k dx^l = (R_{ijkl}^B e^i e^j e^k + R_{kl}^A e^k) dx^l.$$

The operator $R_{kl}^A e^k dx^l = \frac{i}{\hbar} \text{ad}\{H_{T_{kl}} e^k dx^l\}$ acts on a section $\mathbf{s} \in \Gamma(B, {}^A\mathcal{W}_B)$ in the following way:

$$\frac{i}{\hbar} [H_{T_{kl}} e^k dx^l, \mathbf{s}] = \frac{i}{\hbar} [e^k, \mathbf{s}]_{\circ} H_{T_{kl}} dx^l + \{H_{T_{kl}}, \mathbf{s}\} e^k dx^l$$

where T_{kl} is an element in $\mathfrak{g}$ on M and hence its action on $\mathbb{A}^{\hbar}(F)$ is defined, so it is also defined on sections of the bundle ${}^A\mathcal{W}_B$: $[H_{T_{kl}}, \mathbf{s}]_{\circ} = \{H_{T_{kl}}, \mathbf{s}\} = T_{kl} \mathbf{s}$.

The operator $\frac{i}{\hbar} \text{ad}_{\circ}\{e^k\} H_{T_{kl}} dx^l$ is of degree -1 . This term has to be added to δ , the initial -1 -degree operator, but as soon as it is present it changes all the equations (40) since there is not only δ in degree -1 anymore.

However if we change δ the iteration method can still be applied yielding a solution for the flat connection. In the central extension the curvature of this connection starts from the term of degree -2 : ${}^A\delta^2 = \frac{i}{\hbar} \text{ad}_{\circ}\{(\omega_{kl} + H_{T_{kl}}) dx^k \wedge dx^l\}$. We are looking for the solution in the form ${}^A\delta = \frac{i}{\hbar} A_{ik} e^i dx^k$. Applied twice it must give this central element:

$$\frac{i}{\hbar} (\omega_{kl} + H_{T_{kl}}) dx^k \wedge dx^l = gr_{(-2)} \left[\frac{i}{\hbar} A_{ik} e^i dx^k, \frac{i}{\hbar} A_{jl} e^j dx^l \right].$$

Obviously, $A_{jl} \in \Gamma(B, A)$ should be in the form of some series in ω and H_T :

$$A_{ik} \omega^{ij} A_{jl} = \omega_{kl} + H_{T_{kl}}, \quad \text{or } A_{ik} = \omega_{im} (1_{mk} + 1/2 \omega^{mn} H_{T_{nk}} + \dots) = \omega_{im} \left(\sqrt{1 + \omega^{-1} H_T} \right)_k^m$$

We have used here the skew symmetry of the forms $\omega_{kl} = -\omega_{lk}$ and $H_{T_{kl}} = -H_{T_{lk}}$. This series converges if the ratio of ω and H_T is much bigger than 1 (i.e. size of the fibres is very small in comparison to the base). It means that the fiberwise symplectic form should be much smaller than the one on the base.

This forces us to introduce a small parameter ϵ :

$$(\omega_{kl} + H_{T_{kl}}) dx^k \wedge dx^l \rightarrow (\omega_{kl} + \epsilon^2 H_{T_{kl}}) dx^k \wedge dx^l.$$

Then the (-1) -degree term in the flat connection should be

$${}^A\delta = \frac{i}{\hbar} \omega_{km} \left(\sqrt{1 + \epsilon^2 \omega^{-1} H_T} \right)_l^m dx^l \text{ad}_{\circ}\{e^k\}.$$

Now the equation for the flat connection D should start with ${}^A\delta$ instead of δ :

$$D = {}^A\delta + \nabla + r.$$

Let X be the following matrix

$$X = \sqrt{1 + \epsilon^2 \omega^{-1} H_T}.$$

Let us write how ${}^A\delta$ acts on a monomial

$$\mathbf{s} = e^{i_1} \otimes \dots \otimes e^{i_m} \otimes f \otimes dx^{j_1} \wedge \dots \wedge dx^{j_n} \in gr_m(\mathcal{W}_B \otimes_{\mathbb{C}[[\hbar]]} A) \otimes \Lambda^n \mathcal{T}^* B, \quad f \in gr_0(A)$$

$$\begin{aligned} {}^A\delta : e^{i_1} \otimes \dots \otimes e^{i_m} \otimes dx^{j_1} \wedge \dots \wedge dx^{j_n} \mapsto \\ \sum_{k=1}^m e^{i_1} \otimes \dots \widehat{e^{i_k}} \dots \otimes e^{i_m} \otimes f \otimes X_l^{i_k} dx^l \wedge dx^{j_1} \wedge \dots \wedge dx^{j_n} \end{aligned}$$

In order to get rid of this correction term X we can “rescale” the bundle $\mathcal{W}_B$ by changing

$$e^k \rightarrow \hat{e}^k = (X^{-1}e)^k \in \mathcal{W}_B \otimes_{\mathbb{C}[[\hbar]]} A$$

After some calculations we arrive at the following commutation relation:

$$[\hat{e}^k, \hat{e}^l]_{\circ} = -i\hbar(\omega + \epsilon^2 H_T)^{kl}. \quad (41)$$

On monomials in $\hat{e}$, ${}^A\delta$ has the same action as δ on monomials in e (Lemma 2.5). We can define ${}^A\delta^{-1}$ and the whole set up in the twisted Weyl bundle becomes a familiar data for Fedosov quantization of a symplectic manifold. This way we reduce the problem to the usual Fedosov quantization and hence our theorem is proved.

We construct a flat connection on the twisted bundle by iterations, r_i being monomials in $\hat{e}$. Uniqueness follows from the condition $\delta^{-1}r = 0$, which gives the same restriction as ${}^A\delta^{-1}r = 0$. The flat sections of this new connection corresponding to the sections of the auxiliary bundle can also be obtained by the recursive procedure.

The only difference we must emphasize is in the characteristic class of deformation. ${}^A\delta^2$ gives that central element which determines the characteristic class of deformation: $\Omega = \omega + \epsilon^2 H_T$. $\square$

5.3. Quantization of the total space.

Here we want to prove that the deformation we got in the previous section is indeed a deformation of functions on M . Since by quantization of functions on M we understand an isomorphism to flat sections of a flat connection of Weyl algebra bundle over M we reformulate the problem as follows. Namely, there is a homomorphism of noncommutative algebras α such that the following diagram commutes:

$$\begin{array}{ccc} C^\infty(M)[[\hbar]] & \longrightarrow & \Gamma(B, \mathcal{F}) \\ Q \downarrow & & \downarrow Q^A \\ \Gamma_{flat}(M, \mathcal{W}_M) & \xrightarrow{\alpha} & \Gamma_{flat}(B, {}^A\mathcal{W}_B) \end{array}$$

First line is an isomorphism of commutative algebras (32). Left vertical arrow is a quantization of a symplectic manifold M (6). Q^A is a lift of the quantization $Q^{\text{fib}}_{\text{bre}}$ (35) to $\Gamma(B, {}^A\mathcal{W}_B)$ followed by quantization in the twisted Weyl algebra bundle (39) from Theorem 5.6.

The Weyl algebra structure is defined on $\Gamma(M, \mathcal{W}_M)$ from the symplectic form (18) on M , and $\Gamma(B, \mathcal{W}_B)$ from the symplectic form ω_B on B . However

Proposition 5.7. *There exists an isomorphism of algebra bundles:*

$$\beta : \Gamma(M, \mathcal{W}_M) \rightarrow \Gamma(B, \mathcal{W}_B \otimes_{\mathbb{C}[[\hbar]]} \Gamma(M/B, \mathcal{W}_{M/B})).$$

Let D^F be a flat connection on $\Gamma(M/B, \mathcal{W}_{M/B})$, such that $\Gamma_{\text{flat}}(M/B, \mathcal{W}_{M/B}) = A$.

Then there exists a flat connection on $\mathcal{W}_M$:

$$D^M = \bar{D} + D^F,$$

such that its flat sections under isomorphism β are mapped to the flat sections of D^A in ${}^A\mathcal{W}_B = \mathcal{W}_B \otimes_{\mathbb{C}[[\hbar]]} A$. It leads to the homomorphism of algebras:

$$\mathbb{A}^{\hbar}(M) \cong \Gamma_{\text{flat}}(B, {}^A\mathcal{W}_B).$$

Proof. Consider the local frames introduced at the end of Section 3: $\{f_\alpha\}$ of vertical tangent bundle $\mathcal{V}M$ and $\{e_i\}$ in $\mathcal{T}B$ at a point $b = \pi(x)$ of B , with dual frames $\{f^\alpha\}$ and $\{e^i\}$.

Using the connection we obtain a local frame on the tangent bundle $\mathcal{T}M = \pi^*\mathcal{T}B \oplus \mathcal{V}M$ at a point x : $\{\pi^*e_i, f_\alpha\}$. Let us denote the dual frame as $\{\bar{e}^i, f^\alpha\}$.

Then the form on M can be written as a block matrix (18), so that in this frame

$$[f^\alpha, f^\beta] = \epsilon^2 \sigma_b^{\alpha\beta}, \quad [\bar{e}^k, \bar{e}^l] = (\pi^*\omega^B + \epsilon^2 H_T)^{kl}.$$

The flat connection D^F defines an isomorphism of algebras of fibrewise deformed functions. The map $\mathcal{W}_M \rightarrow {}^A\mathcal{W}_B : \bar{e} \mapsto \hat{e}$ defines the algebra homomorphism:

$$\Gamma_{\text{fibrewise flat}}(M, \mathcal{W}_M) \cong \Gamma(B, \mathcal{W}_B \otimes_{\mathbb{C}[[\hbar]]} A).$$

Indeed although the Moyal–Vey products are different, the product $\hat{e}$ in ${}^A\mathcal{W}_B$ and $\bar{e}$ in $\mathcal{W}_M$ give the same $*$ -product (see (41)), so it gives a homomorphism of algebras.

The last step is to construct a connection $\bar{D}$ in $\Gamma_{\text{fibrewise flat}}(M, \mathcal{W}_M)$ so that flat sections of D^A (38) correspond to the flat sections of $\bar{D}$. We do it just by writing for $\bar{D}$ the same formula as for D^A but with $\bar{e}$ instead of $\hat{e}$.

The connection $\bar{D}$ together with the Fedosov connection along the fibres D^F gives rise to a single connection D^M on $\mathcal{W}$:

$$D^M = D^F + \bar{D}.$$

Its flat sections $s \in \Gamma(M, \mathcal{W})$ satisfy simultaneously two equations $D^F s = 0$, $\bar{D}s = 0$. So we get a map from the quantization of the total space to the quantization on the base with values in the auxiliary bundle A . $\square$

Interpretation in MM-calculus. In order to see how this homomorphism works we want to introduce calculus similar to the one in [26].

There is some analogy with Riemannian fibrations (see for example [4]). Mazzeo and Melrose ([26]) gave an interpretation of Hodge–Leray spectral sequence from an analytic point of view. In particular they introduced language similar to b -calculus for description of Riemannian fibrations. The idea was to put in a small parameter ϵ , so that all horizontal differential forms had the parameter in some degree. So everything which came from the base was “marked” by this small parameter. This gave the description of terms in the spectral sequence by the coefficients in Taylor decomposition with respect to this small parameter.

Symplectic fibrations provide somewhat similar picture. One even has the parameter naturally coming in the construction of a symplectic form on the total space. Indeed, when the parameter is zero one gets a fibrewise noncommutative product while along the base it is commutative. The $*$ -product then is a bidifferential expression only in vertical coordinates. The term at the first degree of the parameter gives a Poisson bracket in the horizontal direction, it is a first order bidifferential expression in the horizontal direction. If the fibration is trivial these bidifferential expressions are fibrewise constant.

Given a connection (12) on M : $\mathcal{T}M = \mathcal{H}M \oplus \mathcal{V}M$ one implements the splitting into the structure of the product manifold $X = M \times [0, \epsilon_0)$, where $\epsilon_0 \ll 1$ is some fixed small number. (We want it to be small enough so that ϵ involved in the symplectic form (16) is bigger than ϵ_0 .) The product $X = M \times [0, \epsilon)$ has an induced fibration, with leaves F and the base $B \times [0, \epsilon)$. Consider the space $\mathcal{L}$ of smooth vector fields on X which are tangent to the fibres, M , of the product structure and which are also tangent to the fibres of the fibration $M \rightarrow B$, above $M_0 = \{\nu = 0\}$. In local coordinates x^j, ξ^k in M , where the x 's give coordinates in B , the elements of $\mathcal{L}$ are the vector fields of the form

$$\sum_{j=1}^{2p} a(x, \xi, \nu) \nu \partial_{x_j} + \sum_{k=1}^{2q} b(x, \xi, \nu) \partial_{\xi_k}.$$

Consider a vector bundle ${}^\nu \mathcal{T}M$ for which $\mathcal{L}$ is the set of sections

$$\mathcal{L} = C^\infty(X, {}^\nu \mathcal{T}M).$$

There is a natural bundle map $\iota_\nu : {}^\nu \mathcal{T}M \rightarrow \mathcal{T}_X M$, the lift of $\mathcal{T}M$ to X . It is an isomorphism except over M_0 , where its range is equal to $\mathcal{V}M$. It is important to define the dual map

$$\iota^\nu : \mathcal{T}_X^* M \rightarrow {}^\nu \mathcal{T}^* M,$$

which range over M_0 is a subbundle which is naturally isomorphic to the bundle of forms on fibres.

Given a connection (12) on M : $\mathcal{T}M = \mathcal{H}M \oplus \mathcal{V}M$, the restriction of ${}^\nu \mathcal{T}^* M$ to the boundary, M_0 , of X naturally splits

$${}^\nu \mathcal{T}^*_{M_0} M = \nu^{-1} \mathcal{T}^* B \oplus \mathcal{V}^* M, \quad u_0 = \nu^{-1} \pi^* \beta + \iota_\nu(\alpha).$$

The exterior powers of ${}^\nu \mathcal{T}^* M$ also split at the boundary so one can define a new bundle of rescaled differential forms on X . Namely,

$${}^\nu \mathcal{A}_x^k(M) = \sum_{j=0}^k \mathcal{A}_x^j(M/B) \oplus \nu^{-(k-j)} \mathcal{A}_{\pi(x)}^{k-j}(B).$$

The symmetric powers of ${}^\nu \mathcal{T}^* M$ have the following decomposition

$${}^\nu S^k \mathcal{T}^* M = \sum_{j=0}^k S^j(M/B) \oplus \nu^{-(k-j)} S^{k-j} \mathcal{T}^* B.$$

This way one can define the rescaled Weyl algebra bundle. At a point $x \in M$ as before one can introduce a local frame $\{f_k\}$ of vertical tangent bundle $\mathcal{V}M$ corresponding to ∂_{ξ_k} and a local frame $\{e_i\}$ in $\mathcal{T}B$, corresponding to ∂_{x_j} at a point $b = \pi(x)$ of B , with dual frames $\{f^k\}$ and $\{e^i\}$. Using

the connection we obtain a local frame on the tangent bundle $\mathcal{T}M = \pi^*\mathcal{T}B \oplus \mathcal{V}M$ at a point x . The differential on the bundle of forms ${}^\nu\mathcal{A}^k(M)$ is given by

$$d = \nu \frac{dx^j}{\nu} \partial_{x_j} + d\xi^k \partial_{\xi_k},$$

so that $d : {}^\nu\mathcal{A}^k(M) \rightarrow {}^\nu\mathcal{A}^{k+1}(M)$. Similarly we can find a symplectic connection on ${}^\nu\mathcal{T}M$ from the initial connection on $\mathcal{T}M$.

Then ${}^\nu D$, the Fedosov connection on rescaled Weyl algebra bundle ${}^\nu\mathcal{W}_M$ will have the Taylor decomposition into degrees of ν . Since Fedosov connection is flat it should give an equation in each degree of ν . The ν -decomposition of $({}^\nu D)^2$ must give 0 in each degree of ν . The result for quantization can be stated as follows:

1. The quantization of M for $\nu = 0$ is $C^\infty(B, \mathbb{A}^h(M/B))$
2. The term at the first power of ν is the product on the base given by the Poisson bracket with values in the quantization of the fiber.
3. n -th power of ν allows one to write a product on the base with values in the product of the fibre up-to the n -th power in $\hbar$.

The total space of a symplectic fibration $M \rightarrow B$ together with the rescaled symplectic form and a connection preserving it make such an F -manifold which is easily associated with an F -bundle and hence gives a map from quantization of the total space to a quantization of the base with values in the auxiliary bundle of fibrewise quantizations.

6. EXAMPLES OF SYMPLECTIC FIBRATIONS AND THEIR QUANTIZATION.

Fedosov quantization provides a way to construct a $*$ -product on a symplectic manifold. In the previous section we showed how Fedosov quantization works for any symplectic fibration. However, step-by-step calculations become complicated very quickly and explicit formulas are readily available only in a few particular cases. Mostly the results which we are getting are of the type that "under quantization some of the variables behave in a certain way".

For the trivial symplectic fibration, that is for the direct product of two F -manifolds (B, ω^B, ∇^B) and (F, σ, ∇^F) we get the direct product of quantizations:

$$\mathbb{A}^h(B \times F) = \mathbb{A}^h(B) \times_{\mathbb{C}[[\hbar]]} \mathbb{A}^h(F),$$

with the characteristic class $\omega^B + \epsilon\sigma$, where ϵ can be arbitrary nonzero number (since automatically for nondegenerate ω and σ for any nonzero ϵ the sum is nondegenerate in the absence of the curvature term).

An example of a symplectic fibration with fibres being $\mathbb{R}^{2n}$ is considered in [13].

Good source of examples of symplectic fibrations are cotangent bundles to fibre bundles with connections (see again [18]). Indeed, consider a bundle X over B , such that X is an associated bundle to a principal G -bundle P with a fibre $F : X = P \times_G F$. Then G acts by symplectomorphisms on the cotangent bundle $\mathcal{T}^*F$ equipped with the canonical symplectic form $d\alpha^F$. We construct a new bundle $M = \mathcal{T}^*X$ over $\mathcal{T}^*B$, which is the pullback of $P \times_G \mathcal{T}^*F$ to $\mathcal{T}^*B$.

The connection, that is a splitting of $\mathcal{T}M$, is inherited from the splitting of $\mathcal{T}X$. Namely, $\mathcal{H}M$ is defined as the preimage of $\mathcal{H}X$.

The symplectic form on M is the canonical symplectic form on the cotangent bundle. However due to the fibre bundle structure we can split it into two parts – one along the fibres and the other one coming from the base. There is no need to introduce the small parameter since we know that the form is nondegenerate as is.

The algebraic index theorem on a cotangent bundle coincides with the analytical index theorem of Atiyah–Singer on the initial space [29]. Our example shows that this way we may obtain the relation on the index of the fibre, the base, and the total space for any fibre bundle (if the transformation group is a compact Lie group).

Now let us move to other particular examples. Consider the following situation: the associated bundle $M \rightarrow B$ to the principal G -bundle with a fibre being T^*G – the cotangent bundle to G . This example is the inverse of the quantum reduction problem and it was discussed in the recent article [14], which is the generalization of [12].

The article [12] treats a symplectic fibration $M \rightarrow B$ with a symplectic fibre being a cylinder. The fibre can be represented as $\mathbb{C}^* = T^*S^1$ – the cotangent bundle to the circle. Locally a point z in M can be described by coordinates (x, r, θ) , θ being an angle coordinate on the cylinder and r the height, while $x = (x_1, \dots, x_{2n})$ denotes coordinates of the base point $\pi(z)$. Then $M = P \times_{U(1)} \mathbb{C}^*$, where P is a principal $U(1)$ -bundle. The symplectic form can be constructed from Proposition (3.4). Let λ be a connection one-form on P . In this particular case:

$$\lambda : TP \rightarrow \mathfrak{g} = \mathbb{R}.$$

Hamiltonian of λ is a function in r only, it does not involve θ . Quantization on the fibres is the Weyl quantization like in $\mathbb{R}^{2n}$ (see (43)). The symplectic connection is flat and one gets from this fibrewise quantization that fibrewise flat sections are expressions in $(r + f^1), (\theta + f^2)$, where (f^1, f^2) are generators in the Weyl algebra bundle corresponding to $(dr, d\theta)$ in T^*F . The resulting flat connection in the Weyl algebra bundle on M is a series in $(r + f^1)$ (it does not involve the other coordinate $(\theta + f^2)$).

Examples with two-dimensional fibres are provided by any Riemannian surfaces (except a torus) as fibres (see [24]).

As a nontrivial example of symplectic fibrations we consider S^2 -bundles. S^2 -bundle $\pi : M \rightarrow B$ can be considered as an associated bundle to a principal $U(1)$ -bundle, P . We construct the symplectic form on $M = P \times_{U(1)} S^2$. The manifold M is presented as the symplectic reduction of (W, Ω) at a regular value 0 of the moment map. The algebra of the group $U(1)$ is simply $\mathbb{R}$. Let $\mathcal{V}P \subset TP$ be the bundle of vertical tangent vectors. The fibre at a point p being $\mathcal{V}_pP \subset T_pP$. A connection 1-form, $\lambda : TP \rightarrow \mathbb{R}$ determines a horizontal subbundle by

$$\mathcal{H}_p = \{v \in T_pP \mid \lambda_p(v) = 0\}$$

This horizontal subbundle induces an injection: $\iota_\lambda : \mathcal{V}^*P \hookrightarrow T^*P$, namely, a vertical cotangent vector $\xi \in \mathcal{V}_p^*P$ is a linear functional on T_pP which vanishes on the horizontal subspace $\mathcal{H}_pP$. The subbundle $\mathcal{V}^*P$ inherits the standard symplectic form from T^*P – $d\alpha_{\text{can}}$. The manifold (Weinstein universal space)

$$W = \mathcal{V}^*P \times S^2$$

carries a natural symplectic form $\Omega = pr_B^* \omega_B + \iota_\lambda^* d\alpha_{\text{can}} + pr_S^* \sigma$, where $pr_B : W \rightarrow B$ and $pr_S : W \rightarrow S^2$ are the obvious projections and σ is a $U(1)$ -invariant volume form on S^2 . Ω is invariant

under the diagonal action of $U(1)$. $\mathcal{V}^*P$ is equivariantly diffeomorphic to $P \times \mathbb{R}$, so the moment map $\mu : W = P \times \mathbb{R} \times S^2 \rightarrow \mathbb{R}$ is given by

$$\mu(p, \eta, z) = h(z) - \eta$$

where $h : S^2 \rightarrow \mathbb{R}$ is the height function, it is a moment map for the action of $U(1)$ on S^2 by rotating about the vertical axis. η is a projection $P \times \mathbb{R} \rightarrow \mathbb{R}$.

The level set $\mu^{-1}(0)$ can be identified with the manifold $P \times S^2$ by the map which takes the form Ω to $pr_B^* \omega_B - d(H_\lambda) + pr_S^* \sigma$ on $P \times S^2$, where $H(p, z) = h(z)$ is the height function on S^2 . This form is equivariant under the $U(1)$ action, to make sure it is nondegenerate we need to introduce a small number ϵ , so that the form on M becomes

$$\omega = pr_B^* \omega_B - \epsilon^2 \{d(H_\lambda) + pr_S^* \sigma\}.$$

Let us consider local coordinates at a point m in M : (x, ξ, θ) , where $x = (x_1, \dots, x_{2n})$ denotes coordinates of the base point $\pi(m)$ while ξ and θ are cylindrical polar coordinates in the fibre, ξ gives a height function and θ is an angle. Then the symplectic form on the fibre is

$$\sigma = d\xi \wedge d\theta.$$

The vertical vector field $\frac{\partial}{\partial \theta}$ has a Hamiltonian ξ :

$$H_{\frac{\partial}{\partial \theta}} = \xi.$$

The auxiliary bundle A is a bundle of quantized functions on fibres, it is associated to $U(1)$ -bundle P . The connection on A is inherited from a connection 1-form by Proposition(5.2). In coordinates it is as follows:

$$\nabla^A \mathbf{s} = d\mathbf{s} + \lambda \{\xi, \mathbf{s}\},$$

where λ is a local 1-form on the base and $\{\cdot, \cdot\}$ is a fibrewise Poisson bracket, $\mathbf{s}$ some section of A . The curvature of this connection is:

$$R^A = \frac{i}{\hbar} \text{ad} H_T = T\{\xi, \cdot\},$$

where T is a 2-form on the base, the curvature of the connection λ . Since a Hamiltonian of any vector field is a linear function of just one coordinate ξ . A flat connection is as follows:

$$D^A = \check{\delta} + \nabla + r,$$

where $\check{\delta} = \text{ad} \omega_{kl} \hat{e}^k dx^l$ and $\hat{e}^k = (\sqrt{(1 + \epsilon^2 \omega^{-1} T \xi)} e)^k$.

As a result one gets that the flat connection in the case of a sphere bundle does not depend on the cylindrical angle coordinate θ . The characteristic class of the deformation with values in the auxiliary bundle of quantization of the fibres is $\omega^B + \epsilon^2 T \xi$.

APPENDIX A. CALCULATIONS FROM DIAGRAMS.

Fedosov quantization produces the system of recursively defined equations. Fedosov proved that there are no obstructions to solutions. In what follows we are trying to show how to obtain these equations step by step from diagrams, presenting all equations at once. Since there are other situations when one has to solve systems of recursive equations we also hope that our presentation might be useful in some other calculations maybe of completely different origin.


A.1. Fedosov connection.

Let $\mathcal{A}^p(W^i) = \Gamma(M, \Lambda^p T^*M \otimes \mathcal{W}^i)$. Let us represent the action of operators constituting the connection

$$D = -\delta + \nabla + r_1 + r_2 + r_3 + r_4 + \dots$$

by arrows pointing in directions corresponding to their degree.

Namely, $r_k : \mathcal{A}^0(W^i) \rightarrow \mathcal{A}^1(W^{i+k})$ is drawn to go from the point corresponding to the level i in the first column to the point in the second column k rows down: δ goes up one row, ∇ is on the same level, r_1 goes down one level and so on. Same operators act between first and second column.



For curvature to be equal 0 we have to get 0 when applying the connection twice to any element $a \in \Gamma(M, \mathcal{W})$: $D^2a = 0$. In other words: the sum of arrows coming to the second column $\mathcal{A}^2(W^i)$ should be 0 for every i .

Showing that this is true for any element in $\mathcal{A}^0(W^i) = \Gamma(M, \mathcal{W}^i)$ for any i will do.

First two terms δ of degree -1 and ∇ of degree 0 are known, our purpose is to find the other terms recursively.

For every degree i we get an equation on operators:

- Level (-2): $\delta^2 = 0$
- Level (-1): $-\delta, \nabla = 0$
- Level (0): $-\delta, r_1 + \nabla^2 = 0$
- Level (1): $-\delta, r_2 + [\nabla, r_1] = 0$
- Level (2): $-\delta, r_3 + [\nabla, r_2] + \frac{[r_1, r_1]}{2} = 0$
- Level (3): $-\delta, r_4 + [\nabla, r_3] + [r_1, r_2] = 0$

All these equations can be written together:

$$D^2 = (-\delta + \nabla + [r, \cdot])^2 = \nabla^2 - [\delta, r] + r^2 = 0$$

where $r = r_1 + r_2 + \dots$. It is solved recursively: in each degree $k \geq 0$ one gets an equation involving r_i for $i \leq k$.

A general proof of the statement that there are no obstructions in this recursive procedure is given in [11] (as well as in other sources). Here let us just show what happens in the first few equations.

Degree -2. The equation is $\delta^2 = 0$. It is satisfied by the Lemma (2.5).

Degree -1. Next one is $[\delta, \nabla] = 0$. It is true by a simple calculation.

For this equation we need that the connection ∇ is torsion-free.

Degree 0. Here is the first nontrivial calculation. We have to find such r_1 that $-\delta, r_1 = \nabla^2$.

a) Existence. First of all:

$$[\delta, \nabla^2] = [\delta, \nabla]\nabla - \nabla[\delta, \nabla]$$

which is 0 by the previous equation.

There is an operator δ^* which is a homotopy for δ .

$$\delta^*\delta^* = 0, \quad \delta\delta^* + \delta^*\delta = id \quad c \tag{42}$$

This c is a number of y 's and dx 's, for example for a term $y^{i_1} \dots y^{i_p} dx^{j_1} \dots dx^{j_q}$ this number $c = p + q$. Let us put

$$r_1 = \delta^*\nabla^2$$

then indeed:

$$[\delta, r_1] = \delta(r_1) = \delta(\delta^*\nabla^2) = \nabla^2$$

and also $\delta^*r_1 = 0$.

b) Uniqueness.

Let $r'_1 = r_1 + \alpha$, such that $\delta^*\alpha = 0$. Then $\alpha = \delta^*\beta$ for some β . Hence, $\delta\delta^*\beta = 0$, because $\delta(r_1 + \alpha) = \delta r_1$.

From (42) we get that $\beta = \delta^*\delta\beta$ and $\alpha = \delta^*\beta = \delta^*(\delta^*\delta\beta) = 0$

Degree 1. $[\delta, r_2] = [\nabla, r_1]$ gives the equation on the operator r_2 .

a) Existence. Again we show

$$[\delta, [\nabla, r_1]] = [[\delta, \nabla], r_1] - [\nabla, [\delta, r_1]] = 0$$

b) Uniqueness. $r_2 = \delta^*([\nabla, r_1])$ similar to the previous one.

Recursively getting similar equations for r_n one finds the Fedosov connection. Here are first few terms:

$$D = \delta + \nabla + \delta^{-1}\nabla^2 + \delta^{-1}\{\nabla, \delta^{-1}\nabla^2\} + \dots$$

A.2. Flat sections and the $\ast$ -product.

The Fedosov connection is flat and its flat sections are in one-to-one correspondence with series in $\hbar$ with functional coefficients. There is a map:

$$Q : C^\infty(M)[[\hbar]] \rightarrow \mathcal{A}^0(W)$$

Given a series:

$$\mathbf{a} = a_0 + \hbar a_1 + \hbar^2 a_2 + \dots$$

one can find uniquely the corresponding flat section of the Weyl algebra bundle

$$A = \mathbf{a} + A_1 + A_2 + A_3 + \dots$$

so that

$$\delta^{-1}(A - \mathbf{a}) = 0.$$

This last condition makes the operator $Q^{-1} : \mathcal{A}^0(W) \rightarrow C^\infty(M)[[\hbar]]$ very simple, namely it is just an evaluation of $A \in \mathcal{A}^0(W)$ at zero value of coordinates along the fibres W . This condition could be changed for any other condition fixing the zero section in $\mathcal{A}^0(W)$ (see [10]).


Again, the condition of flatness:

$$DA = 0$$

can be represented by the fact that for all i sum of operators r_k which get to $\mathcal{A}^1(W^i)$ must be 0. It again gives a recursive system of equations.

We notice that all r_k kill functions, because r_k acts as adjoint operators and functions are in the center of $\mathcal{A}^0(W)$, so $r_k(a_i) = 0$. Hence first few equations following from the diagram bellow are

1. $\nabla a_0 - \delta A_1 = 0$
2. $\nabla A_1 - \delta(\hbar a_1 + A_2) = 0$
3. $r_1 A_1 + \nabla(\hbar a_1 + A_2) - \delta A_3 = 0$
4. $r_2 A_1 + r_1 A_2 + \nabla A_3 - \delta(\hbar^2 a_2 + A_4) = 0$
5. $r_3 A_1 + r_2 A_2 + r_1 A_3 + \nabla(\hbar^2 a_2 + A_4) - \delta A_5 = 0$



Let dx^i be a local frame in $\mathcal{T}^*M$. Then let the corresponding generators in W be $\{e^i\}$. Then A_i are of the form $A_{k_1 \dots k_i} e^{k_1} \dots e^{k_i}$.

The symplectic connection ∇ locally can be written as:

$$\nabla = dx^j \frac{\partial}{\partial x^j} + \frac{i}{2\hbar} [\Gamma_{jkl} dx^j e^k e^l, \]$$

So for the first terms of the flat section corresponding to $\mathbf{a} = a_0$ we get:

1. $A_1 = \delta^{-1} \nabla \mathbf{a} = \delta^{-1} d\mathbf{a} = \partial_l \mathbf{a} e^l$
2. $A_2 = \delta^{-1} \nabla A_1 = \delta^{-1} \nabla (\partial_l \mathbf{a} e^l) = \{\partial_k \partial_l \mathbf{a} + \Gamma_{kl}^j \partial_j \mathbf{a}\} e^k e^l$

The first few terms in the $*$ -product of two functions $\mathbf{a}, \mathbf{b} \in C^\infty(M)$ are:

$$\mathbf{a} * \mathbf{b} = \mathbf{a}\mathbf{b} - \frac{i\hbar}{2} \omega^{ij} \partial_i \mathbf{a} \partial_j \mathbf{b} - \hbar^2 (\partial_i \partial_j \mathbf{a} + \Gamma_{ij}^l \partial_l \mathbf{a}) \omega^{im} \omega^{jn} (\partial_m \partial_n \mathbf{b} + \Gamma_{mn}^k \partial_k \mathbf{b}) + \dots$$

A.3. Examples of deformation quantization of symplectic manifolds.

The procedure of deformation quantization requires calculations which are not obvious and most of the time do not give nice formulas. However in few cases one can calculate explicitly the $*$ -product for particular manifolds. The first trivial example is the quantization of $\mathbb{R}^{2n}$. Let $\{x^1, \dots, x^{2n}\}$ be a local coordinate system at some point $x \in \mathbb{R}^{2n}$. The Darboux symplectic form in these coordinates

is

$$\omega = dx^i \wedge dx^{i+n}, \quad 1 \leq i \leq n$$

The standard symplectic form and the trivial connection $\nabla = d$ give an algebra of pseudodifferential operators in $\mathbb{R}^{2n}$. Namely, using calculations from the diagrams in the last two sections we get the Fedosov connection to be:

$$D = d - \delta \quad \text{or in coordinates} \quad D = dx^i \left(\frac{\partial}{\partial x^i} - \frac{\partial}{\partial e^i} \right).$$

Flat section of such connection corresponding to a function $\mathbf{a}$ under the quantization map is as follows

$$A = \mathbf{a} + e^i \frac{\partial \mathbf{a}}{\partial x^i} + e^i e^j \frac{\partial^2 \mathbf{a}}{\partial x^i \partial x^j} + \dots$$

We see that it gives a formula for Taylor decomposition of a function $\mathbf{a}$ at a point x . In fact the e^i terms can be considered as jets. Then the $*$ -product of two flat sections is given by the formula (2). It is easy to deduce that for two functions $\mathbf{a}$ and $\mathbf{b}$ the $*$ -product is

$$\begin{aligned} \mathbf{a} * \mathbf{b} &= \exp \left\{ -i\hbar \frac{\partial}{\partial y^k} \frac{\partial}{\partial z^{n+k}} \right\} \mathbf{a}(\mathbf{y}) \mathbf{b}(\mathbf{z})|_{\mathbf{y}=\mathbf{z}=x} \\ &= \mathbf{a}\mathbf{b} - i\hbar \frac{\partial \mathbf{a}}{\partial x^k} \frac{\partial \mathbf{b}}{\partial x^{n+k}} - \frac{\hbar^2}{2} \left(\frac{\partial^2 \mathbf{a}}{\partial x^k \partial x^l} \right) \left(\frac{\partial^2 \mathbf{b}}{\partial x^{n+k} \partial x^{n+l}} \right) + \dots \end{aligned} \quad (43)$$

Let us map $C^\infty(\mathbb{R}^{2n})$ to differential operators on $\mathbb{R}^n$, considered as polynomials on $T^*\mathbb{R}^n$. Then the $*$ -product gives exactly the product of differential symbols.

Remark A.1. The same scheme actually works for any cotangent bundle T^*M with the canonical symplectic form – the quantized algebra of functions on T^*M is isomorphic to the algebra of differential operators on M . This observation leads to various types of index theorems [19], also [17],[27],[28].

Explicit formulas for quantization of Kähler manifolds were given in [20]. Tamarkin ([31]) showed that for symmetric Kähler manifolds the Fedosov connection can be constructed to have only three summands, it has no terms of degree higher than 1, i.e.

$$D = \delta + \nabla + r_1.$$

This gives compact formulas for the $*$ -product on such manifolds as the 2-sphere, any projective space $\mathbb{C}P^n$, and Grassmanians. The case of $\mathbb{C}P^n$ was also considered in ([5]).

REFERENCES

- [1] V.I.Arnol'd. *Mathematical Methods of Classical Mechanics*. Springer Verlag, 1989
- [2] F.Baehn, M.Flato, C.Fronsdal, A.Lichnerowicz, D.Sternheimer. *Deformation theory and quantization, I,II*. Ann.Phys.**111**(1978), 61-151
- [3] N.Berline, E. Getzler, M.Vergne. *Dirac Operators and Index Theorem*, Springer Verlag, 1992
- [4] J.-M.Bismut. *The Atiyah-Singer index theorem for families of Dirac operators: Two heat equation proofs*, Invent.math. **83**(1986), 91-151
- [5] M.Bordemann, M.Brischle, C.Emmrich, S.Waldmann. *Phase Space Reduction for Star-Products: An Explicit Construction for CP^n* Preprint **q-alg/9503004**.
- [6] A.Connes, M.Flato, D. Sternheimer. *Closed star products and cyclic cohomology*, Let. Mat. Phys.**24** (1992), 1-12
- [7] P. Deligne. *Déformations de l'algèbre des fonctions d'une variété symplectique: comparaison entre Fedosov et De Wilde, Lecomte*, Selecta Math,**1**,(4),1995

- [8] M.De Wilde, P.B.A.Lecompte. *Existence of star-products and of formal deformations in Poisson Lie algebra of arbitrary symplectic manifold*, Lett. Math. Phys. **7** (1983), 487–496
- [9] J.Donin. *On the quantization of Poisson brackets*, Preprint **q-alg/9505028**, Int. J. Mod. Phys. A11 (1996) 863.
- [10] C.Emmrich, A.Weinstein. *The Differential Geometry of Fedosov's Quantization*, Preprint **hep-th/9311094**.
- [11] B.Fedosov. *A simple geometric construction of deformation quantization*, J.of Diff.Geom. **40** (1994), 213–238
- [12] B.Fedosov. *Reduction and eigenstates in deformation quantization*, in Advances in partial differnetial equations, Akademie Verlag, Berlin (1994), 277–297
- [13] B.Fedosov. *Deformation Quantization and Index Theory*, Akademie Verlag GmbH, Berlin 1996
- [14] B.Fedosov. *Non-Abelian Reduction in Deformation Quantization*, Preprint 1997
- [15] I.M.Gelfand, D.A.Kazhdan, D.B.Fuks. *The action of infinite dimensional Lie algebras*, Funct. Anal. Appl. **6** (1972), 9–13
- [16] I.Gelfand, V.Retakh, M.Shubin. *Fedosov manifolds*, **dg-ga/9707024**
- [17] E.Getzler. *The Bargmann representation, generalized Dirac operators and the index of pseudodifferential operators on R^n* , Contemporary Math. (Proceedings of conference at Keio University, July 1994.)
- [18] V.Guillemin, E.Lerman, S.Sternberg. *Symplectic fibrations and multiplicity diagrams*, Cambridge University Press, 1995
- [19] S.Gutt. *An explicit $\ast$ -product on the cotangent bundle of a Lie group*, Lett. in Math.Phys. **7** (1983), 249–258
- [20] A.V.Karabegov. *Deformation quantization with separation of variables on a Kähler manifold*, Preprint **hep-th/9508013**
- [21] M.Kontsevich. *Deformation quantization of Poisson manifolds, I*, Preprint **q-alg/9709040**
- [22] P.B.A. Lecompte *Existence des star-produits selon B.Fedosov*, Séminaire **95.005**, Institut de Mathématique Université de Lège, (1995)
- [23] A.Lichnerowicz. *Connexions symplectiques et $\ast$ -produits invariants*, C.R.Acad.Sc.Paris, t.**291** (1980), 413–417
- [24] D.McDuff, D.Salamon. *Introduction to Symplectic Topology*, Clarendon Press. Oxford, 1995
- [25] J.E.Marsden, T.Ratiu, G.Raugel. *Symplectic connections and the linearization of Hamiltonian systems*, Proc. of Royal Soc. of Edinburgh, **117A** (1991), 329–380
- [26] R.Mazzeo, R.Melrose. *The adiabatic limit, Hodge cohomology and Leray's spectral sequence for a fibration*, J.Differential Geometry **21** (1990) 185–213
- [27] R.Nest, B.Tsygan. *Algebraic index theorem*, Comm. Math. Phys. **172** (1995) No.2, 223–262
- [28] R.Nest, B.Tsygan. *Algebraic index theorem for families*, Advances in Math. vol.113, No 2 (1995), 151–205
- [29] R.Nest, B.Tsygan. *Formal versus analytic index theorems*, Intl.Math.Res.Not. 1996, No.11, 557–564 (1996)
- [30] S.Sternberg. *Minimal coupling and the symplectic mechanics of a classical particle in the presence of a Yang-Mills field*, Proc. Natl. Acad. Sci. USA, **74** (1977), 5253–5254
- [31] D.Tamarkin. *Quantization of symmetric Kähler manifolds*, preprint (1996)
- [32] A.Weinstein. *A universal phase space for particles in Yang-Mills fields*, Letters in Mathem. Phys. **2** (1978), 1–13
- [33] P.Xu. *Fedosov $\ast$ products and quantum momentum maps*, I.H.E.S. preprint, IHES/M/96/57
- [34] A.Weinstein. *Deformation quantization [d'après B. Fedosov and others]*, Séminaire BOURBAKI, 46ème année, 1993-94, n° 789 (1994)